

GLUEING SILTING OBJECTS

QUNHUA LIU, JORGE VITÓRIA, DONG YANG

ABSTRACT. Recent results by Keller and Nicolás and by Koenig and Yang have shown bijective correspondences between suitable classes of t-structures and co-t-structures with certain objects of the derived category: silting objects. On the other hand, the techniques of glueing (co-)t-structures along a recollement play an important role in the understanding of derived module categories. Using the above correspondence with silting objects we present explicit constructions of glueing of silting objects, and, furthermore, we answer the question of when is the glued silting tilting.

1. INTRODUCTION

Keller and Vossieck established in [18] a bijection between bounded t-structures and equivalence classes of silting objects for the bounded derived category of modules over the path algebra over a field of a Dynkin quiver. Recently, this bijection has been extended by Keller and Nicols in [17] and by Koenig and Yang in [19] for finite dimensional algebras over a field. Indeed, they show that such a bijection with silting objects exists exactly when we consider bounded t-structures whose hearts are length categories. It turns out that Keller and Vossieck's result is a corollary, since in finite representation type all hearts of bounded t-structures are length categories. Recently the notion of co-t-structures was introduced independently by Bondarko in [9] and Pauksztello in [23], and a new correspondence between silting objects and bounded co-t-structures was proved in [17]. This bijection will be central in our approach.

Silting objects play, thus, a more general role than tilting objects. They describe all hearts which are length categories and these turn out to be precisely those which are module categories over some finite dimensional algebra over a field. This algebra, although not in general derived equivalent to the one we started with, is obtained as the endomorphism algebra of the silting object - just like in the tilting setting. Indeed, it is easy to observe that a silting object is tilting if and only if it lies in the heart of the corresponding t-structure.

Glueing techniques with respect to a recollement, due to Beilinson, Bernstein and Deligne, have been intensively studied in [6] for t-structures and, recently, in [9] for co-t-structures. This leads to the natural question of how to glue silting objects and which silting objects are glued from *smaller* ones. Indeed, recent work by the two first authors has shown that, in the piecewise hereditary case, all bounded t-structures with length heart are glued with respect to a non-trivial recollement [21]. In this setting it is then clear that every silting object can be decomposed by this process into as many pieces as derived simple factors of the algebra (check [4] and [5] and for terminology). It turns out, however, that an answer to the problem of glueing silting objects can be given more easily when the focus is on co-t-structures rather than on t-structures. Our main result is as follows.

The second and third named authors are supported by DFG - SPP Darstellungstheorie 1388. The authors would like to thank David Pauksztello for helpful remarks and thank David Pauksztello and Lutz Hille for a discussion which led to example 4.4.

Theorem (Theorem 3.1) *Let $\mathcal{R}$ be a recollement of a triangulated category $\mathcal{D}$ of the form*

$$\begin{array}{ccc} & \xleftarrow{i^*} & \xleftarrow{j_!} \\ \mathcal{Y} & \xrightarrow[i^!]{i_*} \mathcal{D} & \xrightarrow[j_*]{j^*} \mathcal{X} \\ & \xleftarrow{i^!} & \xleftarrow{j_*} \end{array}$$

Let X and Y be respectively silting objects of $\mathcal{X}$ and $\mathcal{Y}$ and $(X_{\geq 0}, X_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ be respectively the associated co-t-structure in $\mathcal{X}$ and $\mathcal{Y}$. Then the induced co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ is associated with the silting object $Z = i_*Y \oplus K_X$, with K_X defined by the triangle

$$i_*\beta_{\geq 1}i^!j_!X \longrightarrow j_!X \longrightarrow K_X \longrightarrow (i_*\beta_{\geq 1}i^!j_!X)[1],$$

where $\beta_{\geq 1}$ is a (non-functorial) choice of truncation for the co-t-structure $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in $\mathcal{Y}$.

Furthermore, the question of when can we glue derived equivalences, i.e., tilting objects, comes as a particular setting of the general context of glueing silting. Similar constructions of tilting objects have been discussed in [20] and [3]. In particular, we will show that the construction in [3] is a particular case of ours. The following is our main theorem concerning tilting.

Theorem (Theorem 4.5) *Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ by $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$, where R, C and B are finite-dimensional algebras over a field of finite global dimension. Let X and Y be tilting objects of $\mathcal{X}$ and $\mathcal{Y}$, respectively. Then $Z = i_*Y \oplus K_X$ is tilting if and only if the following conditions hold.*

- (a) $\text{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k]) = 0$ for all $k < -1$;
- (b) $\text{Hom}_{\mathcal{Y}}(i^*j_*X, Y[k]) = 0$ for all $k < 0$;
- (c) $\text{Hom}_{\mathcal{Y}}(i^*j_*X, i^*j_*X[k]) = 0$ for all $k < -1$.

This paper is structured as follows. In the next section we discuss some preliminary results on recollements, t-structures, co-t-structures and silting needed for the later sections. In section 3 we show how to glue silting and we use this in section 4 to give necessary and sufficient conditions for the glued silting to be tilting. These conditions are particularly nice in the hereditary case and they are made explicit in section 5. Finally, in section 6, we make a few observations on the glueing of HRS-tilts with view towards a comment on the compatibility of silting mutation with glueing.

2. PRELIMINARIES

Throughout, $\mathbb{K}$ denotes a fixed algebraically closed field and R, B and C finite dimensional $\mathbb{K}$ -algebras. $\mathcal{D}^b(R)$ will denote the bounded derived category of finitely generated right R -modules and $\mathcal{K}^b(\text{proj-}R)$ the bounded homotopy category of finitely generated projective right R -modules. The symbols $\mathcal{X}, \mathcal{Y}$ and $\mathcal{D}$ denote triangulated categories.

2.1. Recollements. A recollement of $\mathcal{D}$ by $\mathcal{X}$ and $\mathcal{Y}$ is a diagram of six triangle functors

$$(2.1) \quad \begin{array}{ccc} & \xleftarrow{i^*} & \xleftarrow{j_!} \\ \mathcal{Y} & \xrightarrow[i^!]{i_*} \mathcal{D} & \xrightarrow[j_*]{j^*} \mathcal{X} \\ & \xleftarrow{i^!} & \xleftarrow{j_*} \end{array}$$

satisfying the following properties:

- (1) (i^*, i_*) , $(i_*, i^!)$, $(j_!, j^*)$, (j^*, j_*) are adjoint pairs;

- (2) $i_*, j_*, j_!$ are full embeddings;
- (3) $i^! \circ j_* = 0$ (and thus also $j^* \circ i_* = 0$ and $i^* \circ j_! = 0$);
- (4) for each $Z \in \mathcal{D}$ there are triangles

$$\begin{aligned} i_* i^! Z &\rightarrow Z \rightarrow j_* j^* Z \rightarrow i_* i^! Z[1] \\ j_! j^* Z &\rightarrow Z \rightarrow i_* i^* Z \rightarrow j_! j^* Z[1]. \end{aligned}$$

The following result allows us to change the sides of a recollement.

Theorem 2.1. ([8, propositions 3.6 and 3.7], [16, proposition 5 and theorem 7])

Suppose $\mathcal{D}$ has a Serre functor S . Then, both $\mathcal{X}$ and $\mathcal{Y}$ have Serre functors ($S_{\mathcal{X}}$ and $S_{\mathcal{Y}}$ respectively) and for any recollement $\mathcal{R}$ of the form (2.1), there are reflected recollements $\mathcal{R}_{\mathcal{U}}$ and $\mathcal{R}_{\mathcal{L}}$ (the upper and the lower reflection, respectively)

$$\mathcal{R}_{\mathcal{U}} : \mathcal{X} \begin{array}{c} \xleftarrow{j^\#} \\ \xrightarrow{j_!} \\ \xleftarrow{j^*} \end{array} \mathcal{D} \begin{array}{c} \xleftarrow{i_\#} \\ \xrightarrow{i^*} \\ \xleftarrow{i_*} \end{array} \mathcal{Y},$$

where $i_\# = S^{-1}i_*S_{\mathcal{Y}}$ and $j^\# = S_{\mathcal{X}}^{-1}j^*S$ are left adjoints of i^* and $j_!$ respectively, and

$$\mathcal{R}_{\mathcal{L}} : \mathcal{X} \begin{array}{c} \xleftarrow{j^*} \\ \xrightarrow{j_+} \\ \xleftarrow{j^+} \end{array} \mathcal{D} \begin{array}{c} \xleftarrow{i_*} \\ \xrightarrow{i^!} \\ \xleftarrow{i_+} \end{array} \mathcal{Y}.$$

where $i_+ = Si_*S_{\mathcal{Y}}^{-1}$ and $j^+ = S_{\mathcal{X}}j^*S^{-1}$ are right adjoints of $i^!$ and j_* respectively.

2.2. t-structures and co-t-structures. A *torsion pair* in $\mathcal{D}$ is a pair of strict full subcategories $(\mathcal{D}', \mathcal{D}'')$, closed under taking direct summands and direct sums, such that

- (1) $\text{Hom}_{\mathcal{D}}(X, Y) = 0$ for all $X \in \mathcal{D}'$ and $Y \in \mathcal{D}''$;
- (2) for each Z in $\mathcal{D}$, there is a triangle $X \rightarrow Z \rightarrow Y \rightarrow X[1]$ where $X \in \mathcal{D}'$ and $Y \in \mathcal{D}''$.

By definition, it is easy to see that

$$\begin{aligned} \mathcal{D}' &= \{X \in \mathcal{D} : \text{Hom}_{\mathcal{D}}(X, Y) = 0, \forall Y \in \mathcal{D}''\} =: {}^\perp \mathcal{D}'', \\ \mathcal{D}'' &= \{Y \in \mathcal{D} : \text{Hom}_{\mathcal{D}}(X, Y) = 0, \forall X \in \mathcal{D}'\} =: \mathcal{D}'^\perp. \end{aligned}$$

A *t-structure* in $\mathcal{D}$ is a pair $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ such that $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 1})$ is a torsion pair and $\mathcal{D}^{\leq 0} \subseteq \mathcal{D}^{\leq 1}$, where $\mathcal{D}^{\leq n} := \mathcal{D}^{\leq 0}[-n]$ and $\mathcal{D}^{\geq n} := \mathcal{D}^{\geq 0}[-n]$ for $n \in \mathbb{Z}$. The subcategory $\mathcal{D}^{\leq 0}$ is called the *t-aisle*, and $\mathcal{D}^{\geq 0}$ is called the *t-coaisle*. The *heart* $\mathcal{D}^{\leq 0} \cap \mathcal{D}^{\geq 0}$ is always an abelian category. We say the heart is *length* if it has only finitely many simple objects and each object in the heart has finite length. A t-structure is said to be *bounded* if it satisfies

$$\bigcup_{n \in \mathbb{Z}} \mathcal{D}^{\geq n} = \mathcal{D} = \bigcup_{n \in \mathbb{Z}} \mathcal{D}^{\leq n}.$$

For all $n \in \mathbb{Z}$, there is a right adjoint to the inclusion of the subcategory $\mathcal{D}^{\leq n}$ in $\mathcal{D}$, called the *truncation at n* and denoted by $\tau^{\leq n}$. Similarly, there is a left adjoint to the inclusion of $\mathcal{D}^{\geq n}$, denoted by $\tau^{\geq n}$. In fact, the triangle above can be written functorially, for any $Z \in \mathcal{D}$ as follows

$$\tau^{\leq 0} Z \rightarrow Z \rightarrow \tau^{\geq 1} Z \rightarrow (\tau^{\leq 0} Z)[1].$$

Moreover, it is possible to define associated cohomological functors

$$H^i : \mathcal{D} \rightarrow \mathcal{D}^{\leq 0} \cap \mathcal{D}^{\geq 0}, \quad H^i(X) = (\tau^{\leq i} \tau^{\geq i} X)[i], \forall X \in \mathcal{D}.$$

If a t-structure is bounded then, for any object $X \in \mathcal{D}$, $H^i(X) = 0$ for all but finitely many $i \in \mathbb{Z}$. A well-known example is the standard t-structure in $\mathcal{D}^b(R)$. The associated cohomological functors are given by the usual cohomology of complexes in $\mathcal{D}^b(R)$.

Similarly, a *co-t-structure* in $\mathcal{D}$ is a pair $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ such that $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq -1})$ is a torsion pair and $\mathcal{D}_{\geq 0} \subseteq \mathcal{D}_{\geq -1}$, where $\mathcal{D}_{\geq n} := \mathcal{D}_{\geq 0}[-n]$ and $\mathcal{D}_{\leq n} := \mathcal{D}_{\leq 0}[-n]$ for $n \in \mathbb{Z}$. The subcategory $\mathcal{D}_{\geq 0}$ is called the *co-t-aisle*, and $\mathcal{D}_{\leq 0}$ is called the *co-t-coaisle*. The *co-heart* $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$ is not, in general, an abelian category, as in the case of t-structures, but it still has the structure of an additive category. A co-t-structure is said to be *bounded* if it satisfies

$$\bigcup_{n \in \mathbb{Z}} \mathcal{D}_{\geq n} = \mathcal{D} = \bigcup_{n \in \mathbb{Z}} \mathcal{D}_{\leq n}.$$

Contrarily to t-structures, the triangle in the definition is not functorial. However, a choice of $X \in \mathcal{D}_{\geq 0}$ (respectively, of $Y \in \mathcal{D}_{\leq -1}$) as above will be denoted by $\beta_{\geq 0}Z$ (respectively, by $\beta_{\leq -1}Z$). A well-known example (see [9], [23]) is the standard co-t-structure in $\mathcal{K}^b(\text{proj-}R)$. Here, the non-functorial choices of $\beta_{\geq 0}Z$ or $\beta_{\leq 0}Z$ for some object $Z \in \mathcal{K}^b(\text{proj-}R)$ are usually given by the *stupid truncations*, consisting in setting suitable entries of the complex Z to be zero.

It is known that t-structures and co-t-structures can be glued (or induced) with respect to a recollement. The same arguments can be used to prove the following result.

Theorem 2.2. *Let $\mathcal{R}$ be a recollement of $\mathcal{D}$ of the form (2.1). Let $(\mathcal{X}', \mathcal{X}'')$ and $(\mathcal{Y}', \mathcal{Y}'')$ be torsion pairs in $\mathcal{X}$ and $\mathcal{Y}$, respectively.*

(1) *There is a glued torsion pair $(\mathcal{D}', \mathcal{D}'')$ in $\mathcal{D}$ defined by:*

$$\mathcal{D}' = \{Z \in \mathcal{D} : j^*Z \in \mathcal{X}', i^*Z \in \mathcal{Y}'\}, \quad \mathcal{D}'' = \{Z \in \mathcal{D} : j^*Z \in \mathcal{X}'', i^!Z \in \mathcal{Y}''\}.$$

(2) ([6, theorem 1.4.10]) *If $(\mathcal{X}', \mathcal{X}''[1])$ and $(\mathcal{Y}', \mathcal{Y}''[1])$ are t-structures, then $(\mathcal{D}', \mathcal{D}''[-1])$ is a t-structure in $\mathcal{D}$;*

(3) ([9, theorem 8.2.3]) *If $(\mathcal{X}', \mathcal{X}''[-1])$ and $(\mathcal{Y}', \mathcal{Y}''[-1])$ are co-t-structures, then $(\mathcal{D}', \mathcal{D}''[1])$ is a co-t-structure in $\mathcal{D}$.*

(4) ([9], [21], [28]) *The glueing of bounded t-structures with length heart is a bounded t-structure with length heart. Also, the glueing of bounded co-t-structures is still bounded.*

If $\mathcal{D}$ admits a Serre functor S , $\mathcal{R}$ is a recollement of $\mathcal{D}$ of the form (2.1) and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ are t-structures in $\mathcal{X}$ and $\mathcal{Y}$ respectively, then there are three naturally associated t-structures in $\mathcal{D}$, obtained by glueing these with respect to $\mathcal{R}$, $\mathcal{R}_U$ and $\mathcal{R}_L$ - we will denote them by $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, $(\mathcal{D}_U^{\leq 0}, \mathcal{D}_U^{\geq 0})$, $(\mathcal{D}_L^{\leq 0}, \mathcal{D}_L^{\geq 0})$, respectively. Similarly, given co-t-structures $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in $\mathcal{X}$ and $\mathcal{Y}$, we get three glued co-t-structures $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$, $(\mathcal{D}_{\geq 0}^U, \mathcal{D}_{\leq 0}^U)$, $(\mathcal{D}_{\geq 0}^L, \mathcal{D}_{\leq 0}^L)$.

2.3. Correspondences with silting objects. An object $M \in \mathcal{D}$ is *silting* if $\text{Hom}_{\mathcal{D}}(M, M[i]) = 0$, for all $i > 0$ and $\mathcal{D}$ is the smallest triangulated subcategory containing M which is closed under direct summands, extensions and shifts. It is *tilting* if, in addition, $\text{Hom}_{\mathcal{D}}(M, M[i]) = 0$, for all $i < 0$. We say that two such objects M and N are equivalent if $\text{add}(M) = \text{add}(N)$. The following result is a consequence of [22, corollary 5.9] (see also [17]).

Theorem 2.3. *There is a bijection between bounded co-t-structures in $\mathcal{D}$ whose co-heart is additively generated by one object and equivalence classes of silting objects of $\mathcal{D}$.*

This bijection can be described as follows.

- Given a co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$, the associated equivalence class is given by a silting object M which is an additive generator of the co-heart, i.e., $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0} = \text{add}(M)$.
- Given a silting object $M \in \mathcal{D}$, the associated co-t-structure is defined as follows: $\mathcal{D}_{\geq 0}$ is the smallest full subcategory closed under direct summands, direct sums and extensions containing $\{M[i] : i \leq 0\}$; similarly, $\mathcal{D}_{\leq 0}$ is the smallest full subcategory closed under direct summands, direct sums and extensions containing $\{M[i] : i \geq 0\}$.

Theorem 2.4. ([17], [19, theorem 7.1]). *In $\mathcal{D}^b(R)$, there are bijections between*

- *the set of bounded t-structures in $\mathcal{D}^b(R)$ with length heart;*
- *the set of equivalence classes of silting objects in $K^b(\text{proj-}R)$;*
- *the set of bounded co-t-structures in $K^b(\text{proj-}R)$.*

Recall that R is of finite global dimension if and only if $\mathcal{D}^b(R)$ admits a Serre functor ([8], [24]). Also, in this case, we have $K^b(\text{proj-}R) \cong \mathcal{D}^b(R)$ and, thus, the correspondence between t-structures and co-t-structures occurs in the same category. Under this assumption, the bijections of the theorem can be made explicit as follows (see [19, section 6] for details).

- Given a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$, we can associate a co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ by taking $\mathcal{D}_{\leq 0} = \mathcal{D}^{\leq 0}$ and $\mathcal{D}_{\geq 0} = {}^\perp \mathcal{D}^{\leq -1}$; In the language of [9], this is the *left adjacent co-t-structure* to the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$.
- Given a co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ in $\mathcal{D}^b(R)$, we can associate a t-structure by taking $\mathcal{D}^{\leq 0} = \mathcal{D}_{\leq 0}$ and $\mathcal{D}^{\geq 0} = \mathcal{D}_{\geq 0}^\perp$; In the language of [9], this is the *right adjacent t-structure* to the co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$.
- Given a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$, the associated silting object is the direct sum of the indecomposable Ext-projective objects in the aisle, i.e., the objects X of $\mathcal{D}^{\leq 0}$ such that $\text{Hom}_{\mathcal{D}^b(R)}(X, Y[1]) = 0$ for all $Y \in \mathcal{D}^{\leq 0}$.
- Given a silting object M in $\mathcal{D}^b(R)$, the associated t-structure is defined as follows

$$\mathcal{D}^{\leq 0} = \left\{ Z \in \mathcal{D}^b(R) : \text{Hom}_{\mathcal{D}^b(R)}(M, Z[i]) = 0, \forall i > 0 \right\}$$

$$\mathcal{D}^{\geq 0} = \left\{ Z \in \mathcal{D}^b(R) : \text{Hom}_{\mathcal{D}^b(R)}(M, Z[i]) = 0, \forall i < 0 \right\}.$$

Remark 2.5. If R is of finite global dimension, as before, then the co-t-coaisle $\mathcal{D}_{\leq 0}$ associated with a silting object M in $\mathcal{D}^b(R)$ coincides with the aisle $\mathcal{D}^{\leq 0}$ associated with the same silting object. In other words, the co-t-structure associated to M is left adjacent to the t-structure associated to M .

Remark 2.6. The structures glued from adjacent t-structures and co-t-structures are not directly related by adjacency, as observed by Bondarko [9, remark 8.2.4.4]. Suppose, however, that $\mathcal{D}$ has a Serre functor and that $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ (respectively, $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$) is a left adjacent co-t-structure to the t-structure $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ (respectively, $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$). Using the descriptions provided in theorem 2.2 and the functors of theorem 2.1 for a fixed recollement $\mathcal{R}$, we have

$$\mathcal{D}_{\leq 0}^U = \mathcal{D}^{\leq 0} = \{ Z \in \mathcal{D} : j^*Z \in \mathcal{X}^{\leq 0} = \mathcal{X}_{\leq 0}, i^*Z \in \mathcal{Y}^{\leq 0} = \mathcal{Y}_{\leq 0} \}.$$

This means that the co-structure $(\mathcal{D}_{\geq 0}^U, \mathcal{D}_{\leq 0}^U)$ glued by the upper reflected recollement $\mathcal{R}_U$ is left adjacent to the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ glued by the recollement $\mathcal{R}$. Similarly, the t-structure $(\mathcal{D}_L^{\leq 0}, \mathcal{D}_L^{\geq 0})$ glued by $\mathcal{R}_L$ is right adjacent to the co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ glued by $\mathcal{R}$.

Remark 2.7. A tilting object T in $\mathcal{K}^b(\text{proj-}R)$ yields equivalences between $\mathcal{D}^b(R)$ and $\mathcal{D}^b(\text{End}(T))$ and between $\mathcal{K}^b(\text{proj-}R)$ and $\mathcal{K}^b(\text{proj-End}(T))$ (see [25, theorem 6.4]). Under these equivalences, the t-structure and the co-t-structure associated to T in $\mathcal{D}^b(R)$ correspond to the standard ones in $\mathcal{D}^b(\text{End}(T))$ and $K^b(\text{proj-End}(T))$, respectively.

2.4. HRS-tilts and silting mutation. There are mutation operations on both t-structures and silting objects. Recall that the definition of a torsion pair in an abelian category is analogous to that of a torsion pair in a triangulated category, replacing the triangle axiom by a short exact sequence.

Theorem 2.8. ([15, proposition 2.1], [10, proposition 2.5]). *Let $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ be a bounded t-structure in a triangulated category $\mathcal{D}$ with heart $\mathcal{A}$ and associated cohomology functors H^i , $i \in \mathbb{Z}$. Suppose that $(\mathcal{T}, \mathcal{F})$ is a torsion pair in $\mathcal{A}$. Then $(\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$ is a t-structure in $\mathcal{D}$, where*

$$\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0} = \{E \in \mathcal{D} : H^i(E) = 0, \forall i > 0, H^0(E) \in \mathcal{T}\}$$

$$\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0} = \{E \in \mathcal{D} : H^i(E) = 0, \forall i < -1, H^{-1}(E) \in \mathcal{F}\}.$$

The t-structure $(\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$ is called *the HRS-tilt* of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to $(\mathcal{T}, \mathcal{F})$. The right mutation of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to a simple object S in its heart, $\mathcal{A}$, is defined by

$$\mu_S^+(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0}) = (\mathcal{D}_{+, S}^{\leq 0}, \mathcal{D}_{+, S}^{\geq 0}) := (\mathcal{D}_{(\perp S, \text{add}(S))}^{\leq 0}, \mathcal{D}_{(\perp S, \text{add}(S))}^{\geq 0})$$

and its left mutation as

$$\mu_S^-(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0}) = (\mathcal{D}_{-, S}^{\leq 0}, \mathcal{D}_{-, S}^{\geq 0}) := (\mathcal{D}_{(\text{add}(S), S^\perp)}^{\leq 0}, \mathcal{D}_{(\text{add}(S), S^\perp)}^{\geq 0})$$

On the other hand, silting mutation was introduced and studied by Buan, Reiten and Thomas in [11] and, independently, by Aihara and Iyama in [1]. We recall its definition.

Definition 2.9. Let $M = X \oplus Y$ be a silting object in $\mathcal{D}^b(A)$. The *right mutation of M at X* , denoted by $\mu_X^+(M)$ is constructed as follows. Let $X \rightarrow L$ be a left $\text{add}(Y)$ -approximation of X and take its cone, say $\tilde{X}$. Then $\mu_X^+(M) = \tilde{X} \oplus Y$. Similarly, the *left mutation of M at X* is constructed by taking the cone, say $\bar{X}$, of a right $\text{add}(Y)$ -approximation of X , $K \rightarrow X$ and taking $\mu_X^-(M) = \bar{X} \oplus Y$. We say that a mutation is *irreducible* if X is indecomposable.

Theorem 2.10. ([19, theorem 8.11]). *Irreducible silting mutations are compatible with the bijections established in theorem 2.4. More precisely, if $S_1, \dots, S_n$ are the simple objects in the heart of a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ and $X_1, \dots, X_n$ are indecomposable Ext-projective objects in $\mathcal{D}^{\leq 0}$ such that $\text{Hom}(X_i, S_i) \neq 0$, for all $1 \leq i \leq n$, then the aisles corresponding to the silting objects $\mu_{X_i}^+(M)$ and $\mu_{X_i}^-(M)$ are $\mathcal{D}_{+, S_i}^{\leq 0}$ and $\mathcal{D}_{-, S_i}^{\leq 0}$ respectively.*

The question of whether these mutations are compatible with glueing is discussed in section 6.

3. GLUEING OF SILTING OBJECTS

The results in this section make explicit in a more general setting the bijection established by Aihara and Iyama in [1, theorem 2.36]. More precisely, the theorem below shows how to glue two silting objects of X and $\mathcal{Y}$ into a silting object of $\mathcal{D}$ with respect to a recollement of the form (2.1) and compatibly with the bijection between silting objects and co-t-structures of theorem 2.3.

Theorem 3.1. Let $\mathcal{R}$ be a recollement of $\mathcal{D}$ of the form (2.1). Let X and Y be silting objects corresponding to co-t-structures $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in $\mathcal{X}$ and $\mathcal{Y}$, respectively. Then the induced co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$ is associated with the silting object $Z = i_*Y \oplus K_X$, with K_X defined by the following triangle

$$i_*\beta_{\geq 1}i^!j_!X \longrightarrow j_!X \longrightarrow K_X \longrightarrow (i_*\beta_{\geq 1}i^!j_!X)[1],$$

where $\beta_{\geq 1}$ is a (non-functorial) choice of truncation for the co-t-structure $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in $\mathcal{Y}$.

Proof. First observe that, since i_* is a fully faithful functor, i_*Y is partial silting and, moreover, it is easily checked to lie in the co-heart of the glued co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$.

Let us consider the following (non-functorial) triangle associated with the co-t-structure in $\mathcal{Y}$

$$i_*\beta_{\geq 1}i^!j_!X \longrightarrow i_*i^!j_!X \longrightarrow i_*\beta_{\leq 0}i^!j_!X \longrightarrow (i_*\beta_{\geq 1}i^!j_!X)[1]$$

and the following universal triangle of the recollement (applied to $j_!X$)

$$i_*i^!j_!X \longrightarrow j_!X \longrightarrow j_*X \longrightarrow (i_*i^!j_!X)[1].$$

Hence, we get the following commutative diagram where the rows are triangles

$$\begin{array}{ccccccc} i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & i_*i^!j_!X & \longrightarrow & i_*\beta_{\leq 0}i^!j_!X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & j_!X & \longrightarrow & K_X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ i_*i^!j_!X & \longrightarrow & j_!X & \longrightarrow & j_*X & \longrightarrow & (i_*i^!j_!X)[1] \end{array}$$

By the octahedral axiom, this can be completed to a commutative diagram

$$\begin{array}{ccccccc} i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & i_*i^!j_!X & \longrightarrow & i_*\beta_{\leq 0}i^!j_!X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ i_*\beta_{\geq 1}i^!j_!X & \longrightarrow & j_!X & \longrightarrow & K_X & \longrightarrow & (i_*\beta_{\geq 1}i^!j_!X)[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ i_*i^!j_!X & \longrightarrow & j_!X & \longrightarrow & j_*X & \longrightarrow & (i_*i^!j_!X)[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ i_*\beta_{\leq 0}i^!j_!X & \longrightarrow & K_X & \longrightarrow & j_*X & \longrightarrow & (i_*\beta_{\leq 0}i^!j_!X)[1] \end{array}$$

the rows of which are again triangles. We will now show that K_X lies in the co-heart of $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$. Indeed, it is clear that $i_*\beta_{\geq 1}i^!j_!X \in \mathcal{D}_{\geq 1}$ and, therefore, $(i_*\beta_{\geq 1}i^!j_!X)[1] \in \mathcal{D}_{\geq 0}$. Clearly we also have that $j_!X \in \mathcal{D}_{\geq 0}$ and, thus, the second row of the diagram shows that $K_X \in \mathcal{D}_{\geq 0}$. Similarly, $i_*\beta_{\leq 0}i^!j_!X \in \mathcal{D}_{\leq 0}$ and $j_*X \in \mathcal{D}_{\leq 0}$ imply that $K_X \in \mathcal{D}_{\leq 0}$, proving that it lies in $\mathcal{D}_{\leq 0} \cap \mathcal{D}_{\geq 0}$.

Finally, it is enough to observe that Z generates $\mathcal{D}$. It is clear that i_*Y and $j_!X$ generate $\mathcal{D}$ since X and Y generate $\mathcal{X}$ and $\mathcal{Y}$ respectively and i_* and $j_!$ are fully faithful functors. But, by the second triangle of the diagram, $j_!X$ can be generated by K_X and an object in the image of i_* , which is generated by i_*Y . Therefore Z generates $\mathcal{D}$ and thus it is silting. $\square$

Remark 3.2. Note that the leftmost morphism $i_*\beta_{\geq 1}i^!j_!X \rightarrow j_!X$ of the triangle in theorem 3.1 corresponds to the chosen morphism $\beta_{\geq 1}i^!j_!X \rightarrow i^!j_!X$ via the adjunction morphism

$$\mathrm{Hom}(i_*\beta_{\geq 1}i^!j_!X, j_!X) \xrightarrow{\simeq} \mathrm{Hom}(\beta_{\geq 1}i^!j_!X, i^!j_!X).$$

Remark 3.3. Note that, in the proof of the theorem, Z is not necessarily a basic silting object even if both X and Y are basic. We can, however, obtain a basic silting object in the respective co-heart by ignoring the multiplicities of the direct summands of Z .

Similarly, we can describe a glueing of silting objects that is compatible with the bijection between silting objects and t-structures for $\mathcal{D} = \mathcal{D}^b(R)$, where R has finite global dimension.

Corollary 3.4. *Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be silting objects corresponding to t-structures $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ in $\mathcal{X}$ and $\mathcal{Y}$, respectively. Suppose that R has finite global dimension. Then the glued t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is associated with the silting object $Z = j_!X \oplus K_Y$, with K_Y defined by the triangle*

$$j_!\alpha_{\geq 1}j^*i_{\#}Y \longrightarrow i_{\#}Y \longrightarrow K_Y \longrightarrow (j_!\alpha_{\geq 1}j^*i_{\#}Y)[1],$$

where $\alpha_{\geq 1}$ is a (non-functorial) choice of truncation for the left adjacent co-t-structure of the t-structure $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ in $\mathcal{X}$.

Proof. This follows from the previous theorem by observing, as in remark 2.6 that $\mathcal{D}_{\leq 0}^U = \mathcal{D}^{\leq 0}$ and that the left adjacent co-t-structure of a t-structure corresponds to the same silting object. $\square$

The following definition settles what we will mean by *glueing silting objects*. As shown by theorem 3.1, it is more natural, in this setting, to consider co-t-structures rather than t-structures.

Definition 3.5. Let $\mathcal{D}, \mathcal{X}, \mathcal{Y}$ be triangulated categories and $\mathcal{R}$ a recollement of the form (2.1). We say that a silting object $Z \in \mathcal{D}$ is *glued from $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$ with respect to $\mathcal{R}$* if Z is obtained by the construction of theorem 3.1, i.e., Z corresponds to the co-t-structure glued from the co-t-structures associated to X and Y with respect to $\mathcal{R}$.

The object K_X of theorem 3.1 can be described in a non-constructive way as follows.

Proposition 3.6. *Let $\mathcal{R}$ be a recollement of a triangulated category $\mathcal{D}$ of the form (2.1). Let X and Y be silting objects corresponding to co-t-structures $(\mathcal{X}_{\geq 0}, \mathcal{X}_{\leq 0})$ and $(\mathcal{Y}_{\geq 0}, \mathcal{Y}_{\leq 0})$ in $\mathcal{X}$ and $\mathcal{Y}$, respectively. Let $Z = i_*Y \oplus K_X \in \mathcal{D}$ be the silting object glued from X and Y with respect to $\mathcal{R}$, compatible with the glued co-t-structure $(\mathcal{D}_{\geq 0}, \mathcal{D}_{\leq 0})$. Then the following holds.*

- (1) K_X is a right $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$ -approximation of j_*X ;
- (2) K_X is a left $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$ -approximation of $j_!X$;
- (3) Up to summands in $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$, K_X is uniquely determined by the following conditions:
 - (i) $j^*K_X = X$;
 - (ii) $i^*K_X \in \mathcal{Y}_{\geq 0}$;
 - (iii) $i^!K_X \in \mathcal{Y}_{\leq 0}$.

Proof. To prove (1), let $C \in \mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$ and $f \in \mathrm{Hom}_{\mathcal{D}}(C, j_*X)$. Since $(i_*\beta_{\leq 0}i^!j_!X)[1] \in \mathcal{D}_{\leq -1}$, it is clear that $\mathrm{Hom}_{\mathcal{D}}(C, (i_*\beta_{\leq 0}i^!j_!X)[1]) = 0$ and thus, by using the defining triangle

$$i_*\beta_{\leq 0}i^!j_!X \longrightarrow K_X \longrightarrow j_*X \longrightarrow (i_*\beta_{\leq 0}i^!j_!X)[1],$$

f factors through K_X , proving (1). Analogously, (2) can be proven using the defining triangle

$$i_*\beta_{\geq 1}i^!j_!X \longrightarrow j_!X \longrightarrow K_X \longrightarrow (i_*\beta_{\geq 1}i^!j_!X)[1].$$

In order to prove (3) observe that, for any object L satisfying properties (i), (ii) and (iii), we have a canonical triangle coming from the recollement $\mathcal{R}$

$$i_* i^! L \longrightarrow L \longrightarrow j_* X \longrightarrow i_* i^! L[1]$$

and, since $i_*(\mathcal{Y}_{\leq 0}) \subseteq \mathcal{D}_{\leq 0}$, we have $i_* i^! L[1] \in \mathcal{D}_{\leq -1}$. Therefore, for any object $C \in \mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$, any map from C to $j_* X$ factors through L , proving that L is a right $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$ -approximation of $j_* X$. Hence, by (1), L differs from K_X by a summand in $\mathcal{D}_{\geq 0} \cap \mathcal{D}_{\leq 0}$. $\square$

4. GLUEING OF TILTING OBJECTS

In this section we investigate necessary and sufficient conditions for the glued silting to be tilting. These conditions will be expressed exclusively in terms of the functors of the recollement and of vanishing conditions on X and $\mathcal{Y}$ rather than in $\mathcal{D}$. In this section R has finite global dimension.

Proposition 4.1. *Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1) with $X = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be silting objects of X and $\mathcal{Y}$, respectively. Then $Z = i_* Y \oplus K_X$ is tilting if and only if the following conditions are satisfied*

- (1) Y is tilting;
- (2) $\text{Hom}_{\mathcal{Y}}(Y, i^! j_! X[k]) = 0$ for all $k < 0$;
- (3) $\text{Hom}_{\mathcal{Y}}(i^* j_* X, Y[k]) = 0$ for all $k < 0$;
- (4) $\text{Hom}_X(X, j^+ K_X[k]) = 0$ for all $k < 0$.

Proof. The induced silting object Z corresponds to the co-t-structure glued along $\mathcal{R}$ and to the t-structure glued along $\mathcal{R}_L$ from the respective structures on X and $\mathcal{Y}$ associated with X and Y respectively. The statement that Z is tilting is equivalent to the statement that Z lies in the heart of $(\mathcal{D}_L^{\leq 0}, \mathcal{D}_L^{\geq 0})$ which, by the description of the glued structures, translates into the conditions

- (i) $i^! Z \in \mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$ or, equivalently, $Y, i^! K_X \in \mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$
- (ii) $j^* Z \in X^{\leq 0}$;
- (iii) $j^+ Z \in X^{\geq 0}$.

To examine these conditions we will use the following triangle defining K_X

$$(4.1) \quad i_* \beta_{\leq 0} i^! j_! X \longrightarrow K_X \longrightarrow j_* X \longrightarrow (i_* \beta_{\leq 0} i^! j_! X)[1].$$

Condition (i) is equivalent to Y being tilting (corresponding to condition (1) of the theorem) and $i^! K_X \in \mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$. By applying $i^!$ to the triangle (4.1) we get that $i^! K_X \cong \beta_{\leq 0} i^! j_! X \in \mathcal{Y}_{\leq 0} = \mathcal{Y}^{\leq 0}$. By remark 2.7, $\beta_{\leq 0}$ corresponds to the stupid truncation in $\mathcal{D}^b(\text{End}(Y))$ and, therefore, $i^! K_X \in \mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq 0}$ is equivalent to the condition that $i^! j_! X \in \mathcal{Y}^{\geq 0}$. This can then be translated to the condition (2) of the theorem, by definition of the t-structure associated with a silting object.

Clearly, since j^* maps the co-heart in $\mathcal{D}$ to the co-heart in X , (ii) is automatically satisfied for any silting Z and, thus, this condition is irrelevant.

Condition (iii) is equivalent to $\text{Hom}_X(X, j^+ Z[k]) = \text{Hom}_{\mathcal{D}}(j_* X, Z[k]) = 0$ for all $k \leq 0$. By splitting Z into its summands $i_* Y$ and K_X and applying adjunction we get precisely conditions (3) and (4) of the theorem. $\square$

Remark 4.2. To simplify our notation, we will use the fact (see [6, section 1.4]) that $i^* j_* = i^! j_![1]$.

In many important cases, the left side of the recollement is just given by $\mathcal{D}^b(\mathbb{K})$.

Corollary 4.3. Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $X = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(\mathbb{K})$. Let X be a silting object in $\mathcal{X}$ and $Y = \mathbb{K}$. If $Z = i_*Y \oplus K_X$ is tilting then there are finite dimensional $\mathbb{K}$ -vector spaces X'_{-1}, X'_0 such that $i^*j_*X \cong X'_{-1}[1] \oplus X'_0$.

Proof. In view of remark 4.2, the condition (2) of proposition 4.1 is equivalent to $H^k(i^*j_*X) = 0$ for all $k < -1$, since $Y = B = \mathbb{K}$. Since the Serre functor of $\mathcal{D}^b(\mathbb{K})$ is isomorphic to the identity functor, the condition (3) of proposition 4.1 is equivalent to $\text{Hom}_{\mathcal{Y}}(Y[k], i^*j_*X) = 0$ for all $k < 0$, which is equivalent to $H^k(i^*j_*X) = 0$ for all $k > 0$. To summarise, if Z is a tilting object, then it follows from proposition 4.1 that $H^k(i^*j_*X) = 0$ for $k \neq -1, 0$. We are done. $\square$

Example 4.4. Let R be the $\mathbb{K}$ -algebra given by the quiver with relations

$$\begin{array}{ccc} & & 2 \\ & \nearrow \alpha & \downarrow \beta \\ 1 & & 3 \\ & \nwarrow \delta & \downarrow \gamma \end{array}, \quad \beta\alpha, \alpha\delta, \delta\gamma.$$

The simple module S_1 supported at 1 is a partial tilting module of projective dimension 2. It has a minimal projective resolution over R given by the exact sequence

$$0 \longrightarrow P_2 \xrightarrow{\beta} P_3 \xrightarrow{\delta} P_1 \longrightarrow S_1 \longrightarrow 0.$$

Rickard and Schofield showed in [26] that S_1 cannot be completed to a tilting module over R . We will strengthen this result by showing that S_1 cannot be completed to a tilting object in $\mathcal{D}^b(R)$.

If e is the idempotent $e_2 + e_3$ then, as a right A -module, R/ReR is isomorphic to S_1 , which does not have self-extensions. It, therefore, follows from [12, theorem 3.1] and [13, section 2] that there is a recollement

$$\mathcal{D}^b(R/ReR) \begin{array}{c} \xleftarrow{i^*} \\ \xrightarrow{i_*} \\ \xleftarrow{i^!} \end{array} \mathcal{D}^b(R) \begin{array}{c} \xleftarrow{j^!} \\ \xrightarrow{j^*} \\ \xleftarrow{j_*} \end{array} \mathcal{D}^b(eRe),$$

where

$$(4.2) \quad \begin{aligned} i^* &= -\overset{L}{\otimes}_R R/ReR, & j^! &= -\overset{L}{\otimes}_{eRe} eR, \\ i_* &= -\overset{L}{\otimes}_{R/ReR} R/ReR, & j^* &= -\overset{L}{\otimes}_R Re, \\ i^! &= \text{RHom}_R(R/ReR, -), & j_* &= \text{RHom}_{eRe}(Re, -). \end{aligned}$$

Fix $Y = R/ReR$ and let $X \in \mathcal{D}^b(eRe)$ be a silting object (e.g. $X = eRe$). Then theorem 3.1 yields a completion $Z = i_*Y \oplus K_X$ of the partial silting object $i_*Y = S_1$ into a silting object. However, as we will show, Z is never tilting and, thus, S_1 cannot be completed to a tilting object.

Let T be a basic tilting object of $\mathcal{D}^b(R)$ which contains S_1 as a direct summand. The functor j^* factors through the canonical projection functor $\pi : \mathcal{D}^b(R) \rightarrow \mathcal{D}^b(R)/\text{thick}(S_1)$, as follows

$$\begin{array}{ccc} \mathcal{D}^b(R) & \xrightarrow{j^*} & \mathcal{D}^b(eRe) \\ \pi \searrow & & \nearrow \simeq \\ & \mathcal{D}^b(R)/\text{thick}(S_1) & \end{array}$$

Here $\text{thick}(S_1)$ is the smallest triangulated subcategory of $\mathcal{D}^b(R)$ containing S_1 and it is a silting subcategory of $\mathcal{D}^b(R)$. Thus, by [1, theorem 2.37], the object $X = j^*(T)$ is silting in $\mathcal{D}^b(eRe)$ and T is equivalent to $S_1 \oplus K_X$. Observe that eRe is the path algebra of the Kronecker quiver

$$2 \begin{array}{c} \xrightarrow{\gamma} \\ \xrightarrow{\beta} \end{array} 3 .$$

We draw the component of the AR quiver of $\mathcal{D}^b(eRe)$ containing the preprojective modules

$$\begin{array}{ccccccc} & & I'_3[-1] & & P'_3 & & P'_{3,1} \\ & \nearrow & & \searrow & \nearrow & \searrow & \nearrow \\ \dots & & I'_2[-1] & & P'_2 & & P'_{2,1} & & \dots \end{array}$$

where P'_2 and P'_3 are respectively the indecomposable projective eRe -module at the vertex 2 and 3. Let $\tau = \tau_{eRe}$ denote the Auslander–Reiten translation of $\mathcal{D}^b(eRe)$. Then according to [27, lemma 3.1], there exists p and q such that $\tau^p P'_2[q]$ is a direct summand of K_X . We claim that

$$H^n(i^* j_*(\tau^p P'_2[q])) = \begin{cases} \mathbb{K} & \text{if } n = -q - 1, -q + 1, \\ 0 & \text{otherwise.} \end{cases}$$

As a consequence of corollary 4.3, $S_1 \oplus K_X$ cannot be a tilting object, a contradiction.

The claim follows from a direct computation, which we show only for $P'_{2,1} = \tau^{-1} P'_2$ (it is analogous for others). For $n \in \mathbb{Z}$, we have

$$\begin{aligned} H^n(i^* j_*(P'_{2,1})) &= H^{n+1}(i^! j_!(P'_{2,1})) = H^{n+1}(\text{RHom}_R(R/ReR, j_!(P'_{2,1}))) \\ &= H^{n+1}(\text{RHom}_R(S_1, j_!(P'_{2,1}))) = \text{Hom}_{\mathcal{D}^b(R)}(S_1, j_!(P'_{2,1})[n+1]). \end{aligned}$$

We take a minimal projective resolution of $P'_{2,1}$

$$P'_2 \xrightarrow{\begin{pmatrix} \gamma \\ \beta \end{pmatrix}} \underline{P'_3 \oplus P'_3},$$

where the underlined term is in degree 0. Applying $j_!$ we obtain the following object in $\mathcal{D}^b(R)$:

$$P_2 \xrightarrow{\begin{pmatrix} \gamma \\ \beta \end{pmatrix}} \underline{P_3 \oplus P_3}.$$

The above differential is injective and its cokernel has the following minimal injective resolution

$$\underline{I_2 \oplus I_2 \oplus I_1} \longrightarrow I_3 \oplus I_3 \oplus I_3 \longrightarrow I_1 \longrightarrow I_2 \longrightarrow I_3.$$

Therefore $j_!(P'_{2,1})$ is isomorphic to this complex in $\mathcal{D}^b(R)$ and it follows that

$$\text{Hom}_R(S_1, j_!(P'_{2,1})[n+1]) = \begin{cases} \mathbb{K} & \text{if } n = -1, 1, \\ 0 & \text{otherwise.} \end{cases}$$

In using proposition 4.1 we do not seem to be dealing with computations in $\mathcal{D}$. Still, these occur in the construction of K_X . This can be avoided by introducing the assumption that X is tilting.

Theorem 4.5. Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1) with $X = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be tilting objects of $\mathcal{X}$ and $\mathcal{Y}$, respectively. Then $Z = i_*Y \oplus K_X$ is tilting if and only if the following conditions are satisfied

- (a) $\text{Hom}_{\mathcal{Y}}(Y, i^* j_* X[k]) = 0$ for all $k < -1$;
- (b) $\text{Hom}_{\mathcal{Y}}(i^* j_* X, Y[k]) = 0$ for all $k < 0$;
- (c) $\text{Hom}_{\mathcal{Y}}(i^* j_* X, i^* j_* X[k]) = 0$ for all $k < -1$.

Proof. Condition (a) is obtained from (2) of the previous theorem by remark 4.2 and condition (b) corresponds exactly to condition (3) of the previous theorem. We will show that if X is tilting and (a) and (b) are satisfied then condition (4) of proposition 4.1 is equivalent to (c). Indeed, if X is tilting, $\text{Hom}_{\mathcal{D}}(j_* X, j_* X[k]) = 0$ for all $k \neq 0$ and thus, applying the functor $\text{Hom}_{\mathcal{D}}(j_* X, -)$ to the k -th shift of triangle (4.1), we get $\text{Hom}_{\mathcal{D}}(j_* X, K_X[k]) \cong \text{Hom}_{\mathcal{D}}(j_* X, (i_* \beta_{\leq 0} i^! j_! X)[k])$ for all $k \neq 0, 1$. Since (j_*, j^+) is an adjoint pair, this shows that condition (4) of proposition 4.1 is equivalent to

$$\text{Hom}_{\mathcal{D}}(j_* X, (i_* \beta_{\leq 0} i^! j_! X)[k]) = \text{Hom}_{\mathcal{D}}(i^* j_* X, (\beta_{\leq 0} i^! j_! X)[k]) = 0, \forall k < 0.$$

Applying the functor $\text{Hom}_{\mathcal{Y}}(i^* j_* X, -)$ to the triangle

$$(\beta_{\geq 1} i^! j_! X)[k] \longrightarrow i^! j_! X[k] \longrightarrow (\beta_{\leq 0} i^! j_! X)[k] \longrightarrow (\beta_{\geq 1} i^! j_! X)[k+1]$$

we get that $\text{Hom}_{\mathcal{Y}}(i^* j_* X, \beta_{\leq 0} i^! j_! X[k]) \cong \text{Hom}_{\mathcal{Y}}(i^* j_* X, i^! j_! X[k])$, for all $k < 0$. Indeed, this follows from (b) after recalling that $\beta_{\geq 1} i^! j_! X$ lies in $\mathcal{Y}_{\geq 1}$, which by construction (see 2.3) is suitably generated by $(Y[n])_{n < 0}$ - showing that $\text{Hom}_{\mathcal{Y}}(i^* j_* X, (\beta_{\geq 1} i^! j_! X)[k+1]) = 0$ for all $k < 0$. Using again remark 4.2, we obtain condition (c), thus finishing the proof. $\square$

Before showing applications of this theorem, we discuss the behaviour of the Serre functor with respect to a t-structure corresponding to a tilting object. Recall from [6, proposition 3.1.10] that for any bounded t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$ with heart $\mathcal{A}$, there is a triangle functor $real : \mathcal{D}^b(\mathcal{A}) \rightarrow \mathcal{D}^b(R)$, t-exact for the standard t-structure $(\mathcal{D}_{st}^{\leq 0}, \mathcal{D}_{st}^{\geq 0})$ in $\mathcal{D}^b(\mathcal{A})$ and $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ in $\mathcal{D}^b(R)$, i.e., it is both right t-exact ($real(\mathcal{D}_{st}^{\leq 0}) \subseteq \mathcal{D}^{\leq 0}$) and left t-exact ($real(\mathcal{D}_{st}^{\geq 0}) \subseteq \mathcal{D}^{\geq 0}$).

Lemma 4.6. Let $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ be a bounded t-structure in $\mathcal{D}^b(R)$ with associated basic silting object T . Let S_R be the Serre functor of $\mathcal{D}^b(R)$. Then the following are equivalent:

- (1) T is tilting;
- (2) S_R is right t-exact;
- (3) $S_R(T)$ lies in the heart.

Proof. (1) $\Rightarrow$ (2): Suppose T is tilting. Let $\Gamma = \text{End}_{\mathcal{D}^b(R)}(T)$ and let $(\mathcal{D}_{st}^{\leq 0}, \mathcal{D}_{st}^{\geq 0})$ be the standard t-structure in $\mathcal{D}^b(\Gamma)$. Then, by [2, theorem 6.6], the realisation functor is a t-exact equivalence. Therefore, $real \circ S_{\Gamma} = S_R \circ real$ (since Serre functors are unique) and $real(\mathcal{D}_{st}^{\leq 0}) = \mathcal{D}^{\leq 0}$. Note also that S_{Γ} preserves the standard aisle in $\mathcal{D}^b(\Gamma)$ since it can be realised as a derived tensor product with a bimodule. This is enough to observe that S_R is right t-exact, since

$$S_R(\mathcal{D}^{\leq 0}) = S_R(real(\mathcal{D}_{st}^{\leq 0})) = real(S_{\Gamma}(\mathcal{D}_{st}^{\leq 0})) \subseteq real(\mathcal{D}_{st}^{\leq 0}) = \mathcal{D}^{\leq 0}.$$

(2) $\Rightarrow$ (3): For any $K \in \mathcal{D}^{\leq -1}$, $\text{Hom}_{\mathcal{D}^b(R)}(K, S_R T) = \text{Hom}_{\mathcal{D}^b(R)}(T, K) = 0$, by definition of the t-structure associated to T (see subsection 2.3). This shows that $S_R T \in \mathcal{D}^{\geq 0}$. If S_R is right t-exact, then $S_R(T) \in \mathcal{D}^{\leq 0}$ and, thus, $S_R(T)$ lies in the heart.

(3) $\Rightarrow$ (1): Suppose $S_R(T)$ lies in the heart. By the definition of the t-structure associated to T , it is clear that $\text{Hom}(T, T[k]) = \text{Hom}(T, S_R(T)[-k]) = 0$ for all $k \neq 0$. Therefore T is tilting. $\square$

Proposition 4.7. *Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1) with $X = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be tilting objects of $\mathcal{X}$ and $\mathcal{Y}$, respectively. If i^*j_*X is an element of $\mathcal{D}^b(B)$ such that $\text{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k])$ is zero except for two consecutive values of $k \in \mathbb{Z}$, then the family of silting objects $(Z_n = i_*Y \oplus K_{X[n]})_{n \in \mathbb{Z}}$ contains at least one tilting object.*

Proof. Let $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ be the t-structure associated with Y . By definition of the t-structure associated to Y , $\text{Hom}_{\mathcal{Y}}(Y, i^*j_*X[k])$ is nonzero exactly for two consecutive values of $k \in \mathbb{Z}$ if and only if there is $a \in \mathbb{Z}$ such that $i^*j_*X \in \mathcal{Y}^{\geq a} \cap \mathcal{Y}^{\leq a+1}$. Since Y is tilting, by lemma 4.6 we have that $S_{\mathcal{Y}}i^*j_*X \in \mathcal{Y}^{\leq a+1}$. In theorem 4.5, condition (a) can be reformulated as $i^*j_*X \in \mathcal{Y}^{\geq -1}$ and condition (b) as $S_{\mathcal{Y}}i^*j_*X \in \mathcal{Y}^{\leq 0}$. Note that, once condition (c) is satisfied for some $X[n]$ it is satisfied for all its shifts. By theorem 4.5, Z_n is tilting if and only if

- (a) $i^*j_*X[n] \in \mathcal{Y}^{\geq -1}$, which means that $a - n \geq -1$ or, equivalently, $n \leq a + 1$;
- (b) $S_{\mathcal{Y}}i^*j_*X[n] \in \mathcal{Y}^{\leq 0}$ which is guaranteed if $a + 1 - n \leq 0$ or, equivalently, $n \geq a + 1$
- (c) $\text{Hom}_{\mathcal{Y}}(i^*j_*X, i^*j_*X[k]) = 0$ for all $k < -1$.

Therefore, take $n = a + 1$ and we only need to check condition (c). It, however, is automatic from our assumption that i^*j_*X has cohomologies with respect to $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ concentrated in exactly two consecutive degrees, thus finishing the proof. $\square$

Remark 4.8. In the family $(Z_n)_{n \in \mathbb{Z}}$ there may be more than one tilting object. Let a be the maximal integer such that $i^*j_*X \in \mathcal{Y}^{\geq a}$ and b be the minimal integer such that $S_{\mathcal{Y}}i^*j_*X \in \mathcal{Y}^{\leq b}$. Then Z_n is tilting if and only if $b \leq n \leq a + 1$.

From the proof above, we see that, for X and Y tilting, whenever i^*j_*X lies in $\mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq -1}$ (i.e., the cohomologies of i^*j_*X with respect to Y lie in degrees -1 and 0), the induced silting $Z = i_*Y \oplus K_X$ is tilting. Since $i^*j_* = i^!j_![1]$, the condition is equivalent to that $i^!j_!X \in \mathcal{Y}^{\leq 1} \cap \mathcal{Y}^{\geq 0}$ which, in its turn, is equivalent to $\text{Hom}_{\mathcal{Y}}(Y, i^!j_!X[k]) \cong \text{Hom}_{\mathcal{D}}(i_*Y, j_!X[k]) = 0$ whenever $k \neq 0, 1$. Under this assumption, we can apply the construction from [3] (theorem 2.5) to the pair $T_1 = j_!X$ and $T_2 = i_*Y$. A similar construction was studied in [20] in the setting of triangular matrix rings.

We briefly recall the construction in [3]. Since $(j_!, j^*)$ is an adjoint pair in the recollement (of the form (2.1)), $\text{Hom}_{\mathcal{D}}(j_!X, i_*Y[k]) \cong \text{Hom}_{\mathcal{X}}(X, j^*i_*Y[k]) = 0$ for all $k \in \mathbb{Z}$. Let m be the dimension of $\text{Hom}_{\mathcal{D}}(i_*Y, j_!X[1])$ and take a basis $\alpha_1, \dots, \alpha_m : i_*Y \rightarrow j_!X[1]$. Consider the universal maps

$$\begin{aligned} \alpha &= (\alpha_1, \dots, \alpha_m)^{tr} : i_*Y^{\oplus m} \rightarrow j_!X[1], \\ \beta &= (\alpha_1, \dots, \alpha_m) : i_*Y \rightarrow j_!X[1]^{\oplus m}. \end{aligned}$$

The map α is left-universal, i.e., that the induced map

$$\text{Hom}(i_*Y, \alpha) : \text{Hom}_{\mathcal{D}}(i_*Y, i_*Y^{\oplus m}) \rightarrow \text{Hom}_{\mathcal{D}}(i_*Y, j_!X[1])$$

is surjective. Similarly, the map β is right-universal, i.e., the induced map

$$\text{Hom}(\beta, j_!X[1]) : \text{Hom}_{\mathcal{D}}(j_!X[1]^{\oplus m}, j_!X[1]) \rightarrow \text{Hom}_{\mathcal{D}}(i_*Y, j_!X[1])$$

is surjective. Consider the triangles determined by α and β :

$$\begin{aligned} T_{\alpha} &: j_!X \rightarrow C_1 \rightarrow i_*Y^{\oplus m} \xrightarrow{\alpha} j_!X[1], \\ T_{\beta} &: j_!X^{\oplus m} \rightarrow C_2 \rightarrow i_*Y \xrightarrow{\beta} j_!X[1]^{\oplus m}. \end{aligned}$$

Theorem 2.5 in [3] asserts that $i_*Y \oplus C_1$ and $j_!X \oplus C_2$ are tilting object in $\mathcal{D}$.

The next proposition proves that the tilting object $i_*Y \oplus C_1$ is precisely the silting (indeed tilting) glued from $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$ with respect to the recollement $\mathcal{R}$, and $j_!X \oplus C_2$ is the silting (indeed tilting) glued from $X \in \mathcal{X}$ and $Y \in \mathcal{Y}$ with respect to the upper reflection $\mathcal{R}_U$. As a consequence, under the condition of proposition 4.5, we are not only able to glue tilting objects in the original recollement $\mathcal{R}$, but also in the upper reflection $\mathcal{R}_U$.

Proposition 4.9. *Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X and Y be tilting objects of $\mathcal{X}$ and $\mathcal{Y}$, respectively, such that $\text{Hom}_{\mathcal{Y}}(Y, i^* j_* X[k]) = 0$ for all $k \neq 0, -1$. Then, up to multiplicity, the tilting object $i_*Y \oplus C_1$ coincides with $Z = i_*Y \oplus K_X$ of theorem 3.1, and the tilting object $j_!X \oplus C_2$ coincides with $Z_U = j_!X \oplus K_Y$ of corollary 3.4.*

Proof. By hypothesis, $i^* j_* X \in \mathcal{Y}^{\leq 0} \cap \mathcal{Y}^{\geq -1}$, for the t-structure $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ associated with Y . We will prove that C_1 satisfies the conditions (i) – (iii) for K_X in proposition 3.6 (3). It follows then $i_*Y \oplus C_1$ lies in the glued co-heart of $\mathcal{D}$ and it generates $i_*Y \oplus j_!X$, thus generating the whole of $\mathcal{D}$. Therefore, up to multiplicities, it coincides with the glued silting $Z = i_*Y \oplus K_X$.

For condition (i), we apply j^* to the defining triangle T_α of C_1 and obtain that $j^*C_1 = j^*j_!X = X$, since $j^*i_* = 0$. For condition (ii), we apply i^* to T_α and obtain that $i^*C_1 = i^*i_*Y^{\oplus m} = Y^{\oplus m} \in \mathcal{Y}_{\geq 0}$, since $i^*j_! = 0$. For condition (iii) we apply $i^!$ to T_α and obtain a triangle

$$i^!j_!X \rightarrow i^!C_1 \rightarrow i^!i_*Y^{\oplus m} \xrightarrow{i^!\alpha} i^!j_!X[1].$$

By applying $\text{Hom}_{\mathcal{Y}}(Y, -)$ to this triangle, we obtain a long exact sequence

$$\dots \rightarrow \text{Hom}_{\mathcal{Y}}(Y, i^!i_*Y^{\oplus m}) \xrightarrow{(i^!\alpha)_*} \text{Hom}_{\mathcal{Y}}(Y, i^!j_!X[1]) \rightarrow \text{Hom}_{\mathcal{Y}}(Y, i^!C_1[1]) \rightarrow \text{Hom}_{\mathcal{Y}}(Y, i^!i_*Y[1]^{\oplus m}) \rightarrow \dots$$

For $k > 0$, we have $\text{Hom}_{\mathcal{Y}}(Y, i^!i_*Y[k]^{\oplus m}) \simeq \text{Hom}_{\mathcal{Y}}(Y, Y[k]^{\oplus m}) = 0$ (since, by assumption, Y is tilting) and for $k > 1$, $\text{Hom}_{\mathcal{Y}}(Y, i^!j_!X[k]) = 0$ (by our assumption on $i^*j_*X = i^!j_!X[1]$). Hence, $\text{Hom}_{\mathcal{Y}}(Y, i^!C_1[k]) = 0$, for all $k > 1$. By adjunction there is a commutative diagram

$$\begin{array}{ccc} (i^!\alpha)_* : & \text{Hom}_{\mathcal{Y}}(Y, i^!i_*Y^{\oplus m}) & \longrightarrow \text{Hom}_{\mathcal{Y}}(Y, i^!j_!X[1]) \\ & \simeq \uparrow & \simeq \uparrow \\ \alpha_* : & \text{Hom}_{\mathcal{D}}(i_*Y, i_*Y^{\oplus m}) & \longrightarrow \text{Hom}_{\mathcal{D}}(i_*Y, j_!X[1]). \end{array}$$

Since, by construction, α is left-universal, i.e., α_* is surjective, it follows that $(i^!\alpha)_*$ is surjective. Hence, we get $\text{Hom}_{\mathcal{Y}}(Y, i^!C_1[1]) = 0$ and, thus, $i^!C_1 \in \mathcal{Y}^{\leq 0} = \mathcal{Y}_{\leq 0}$. The proof for statement about $j_!X \oplus C_2$ is analogous. $\square$

5. THE HEREDITARY CASE

In this section, assuming that R has finite global dimension, we apply theorem 4.5 when B is a hereditary algebra. The conditions for the glued silting to be tilting are then easier to handle.

Proposition 5.1. *Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $\mathcal{X} = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(B)$. Let X be a tilting object in $\mathcal{X}$ and $Y = B$. Assume that B is hereditary. Then $Z = i_*Y \oplus K_X$ is tilting if and only if there are finitely generated B -modules X'_{-1} , X'_0 and X'_1 such that*

- i^*j_*X is isomorphic to $X'_{-1}[1] \oplus X'_0 \oplus X'_1[-1]$;
- X'_1 is either zero or not projective;
- $\text{Hom}_B(X'_1, X'_{-1}) = 0$.

Proof. We analyse the conditions of theorem 4.5 when B is hereditary and $Y = B$. Since B is hereditary, we may assume that $i^* j_* X = \bigoplus_{n \in \mathbb{Z}} X'_n[-n]$, with X'_n a finitely generated right B -module for all n in $\mathbb{Z}$. In this setting, $\text{Hom}_{\mathcal{Y}}(Y, i^* j_* X[k]) = 0$, for all $k < -1$ is equivalent to

$$\text{Hom}_{\mathcal{D}^b(B)}(B, \bigoplus_{n \in \mathbb{Z}} X'_n[-n+k]) = 0 \text{ for all } k < -1$$

which happens if and only if $X'_k = 0$, for all $k < -1$. On the other hand, the condition

$$\text{Hom}_{\mathcal{Y}}(i^* j_* X, Y[k]) = \text{Hom}_{\mathcal{D}^b(B)}(i^* j_* X, B[k]) = 0, \text{ for all } k < 0$$

can be reformulated, using the Serre functor S_B in $\mathcal{D}^b(B)$, by

$$\text{Hom}_{\mathcal{D}^b(B)}(B, S_B(i^* j_* X)[k]) = 0, \text{ for all } k > 0.$$

It is well known that $S_B = [1] \circ \tau_B$, where τ_B is the Auslander-Reiten translation in $\mathcal{D}^b(B)$ and, thus, the statement above is equivalent to

$$\text{Hom}_{\mathcal{D}^b(B)}(B, \bigoplus_{n \in \mathbb{Z}} \tau_B(X'_n)[-n+k+1]) = 0, \forall k > 0,$$

i.e., $X'_k = 0$, for all $k > 1$, and X'_1 is not a projective B -module. So, (a) and (b) of theorem 4.5 hold if and only if $i^* j_* X = X'_{-1}[1] \oplus X'_0 \oplus X'_1[-1]$, with X'_1 not projective as a B -module. Assuming this, we can easily unfold the last condition of theorem 4.5 as follows

$$\text{Hom}_{\mathcal{Y}}(i^* j_* X, i^* j_* X[k]) = 0, \forall k < -1 \Leftrightarrow \text{Hom}_{\mathcal{D}^b(B)}(X'_1, X'_{-1}) = 0,$$

thus finishing the proof. $\square$

As a nice corollary, we obtain a necessary and sufficient condition for the case $\mathcal{Y} = \mathcal{D}^b(\mathbb{K})$.

Corollary 5.2. *Let $\mathcal{R}$ be a recollement of $\mathcal{D} = \mathcal{D}^b(R)$ of the form (2.1), with $X = \mathcal{D}^b(C)$ and $\mathcal{Y} = \mathcal{D}^b(\mathbb{K})$. Let X be a tilting object in $\mathcal{X}$ and $Y = \mathbb{K}$. Then $Z = i_* Y \oplus K_X$ is tilting if and only if there are finite dimensional $\mathbb{K}$ -vector spaces X'_{-1}, X'_0 such that $i^* j_* X \cong X'_{-1}[1] \oplus X'_0$.*

Example 5.3. Let R be the path algebra over $\mathbb{K}$ of the quiver $1 \longrightarrow 2$, of type A_2 . Let $e = e_1$ be the trivial path at the vertex 1. Consider the following standard recollement

$$\begin{array}{ccccc} & \xleftarrow{i^*} & & \xleftarrow{j_!} & \\ \mathcal{D}^b(R/ReR) & \xrightarrow{i_*} & \mathcal{D}^b(R) & \xrightarrow{j^*} & \mathcal{D}^b(eRe) \\ & \xleftarrow{i^!} & & \xleftarrow{j_*} & \end{array}$$

where the six functors are defined as in (4.2). Let D denote the functor $\text{Hom}_{\mathbb{K}}(-, \mathbb{K})$. It is easy to observe that:

- (1) as algebras both R/ReR and eRe are isomorphic to k ;
- (2) as an R -module eR is the simple module S_1 , supported at the vertex 1, and has an injective resolution $(D(Re) \rightarrow D(R(1-e)))$;
- (3) as an R -module R/ReR is the simple module S_2 supported at the vertex 2.

Fix $Y = R/ReR$ and $X = eRe$. Then, in $\mathcal{D}^b(R)$, we have

$$\begin{aligned} i^! j_! X &= i^!(eR) = \text{RHom}_R(R/ReR, eR) \\ &= (\text{Hom}_R(R/ReR, D(Re)) \rightarrow \text{Hom}_R(R/ReR, D(R(1-e)))) \\ &= R/ReR[-1]. \end{aligned}$$

By corollary 5.2, there are exactly two tilting objects in $\{Z_n = i_*Y \oplus K_{X[n]} : n \in \mathbb{Z}\}$. We construct this family using theorem 3.1. For $n \in \mathbb{Z}$, we can choose the map $\beta_{\geq 1}(i^! j_! X[n]) \rightarrow i^! j_! X[n]$ to be

$$\begin{cases} R/ReR[n-1] \xrightarrow{id} R/ReR[n-1] & \text{if } n \leq 0, \\ 0 \rightarrow R/ReR[n-1] & \text{if } n > 0. \end{cases}$$

By adjunction we have a bijection

$$\text{Hom}(i_* \beta_{\geq 1}(i^! j_! X[n]), j_! X[n]) \xrightarrow{\cong} \text{Hom}(\beta_{\geq 1}(i^! j_! X[n]), i^! j_! X[n]).$$

If $n \geq 0$, this adjunction morphism is

$$\text{Hom}_{\mathcal{D}^b(R)}(S_2[n-1], S_1[n]) \xrightarrow{\cong} \text{Hom}_{\mathcal{D}^b(R)}(R/ReR[n-1], R/ReR[n-1]).$$

Both vector spaces are 1-dimensional over $\mathbb{K}$. Let f be the preimage of $id_{R/ReR[n-1]}$. Then the cone of $f[-n]$ is $(1-e)R$, the projective cover P_2 of S_2 . If $n < 0$, the above adjunction morphism is

$$\text{Hom}_{\mathcal{D}^b(R)}(0, S_1[n]) \xrightarrow{\cong} \text{Hom}_{\mathcal{D}^b(R)}(0, R/ReR[n-1]).$$

Thus a morphism $i_* \beta_{\geq 1}(i^! j_! X[n]) \rightarrow j_! X[n]$ as in theorem 3.1 is (see remark 3.2)

$$\begin{cases} S_2[n-1] \xrightarrow{f} S_1[n] & \text{if } n \leq 0, \\ 0 \rightarrow S_1[n] & \text{if } n > 0. \end{cases}$$

Therefore

$$K_{X[n]} = \begin{cases} P_2[n] & \text{if } n \leq 0, \\ S_1[n] & \text{if } n > 0, \end{cases} \quad \text{and} \quad Z_n = \begin{cases} S_2 \oplus P_2[n] & \text{if } n \leq 0, \\ S_2 \oplus S_1[n] & \text{if } n > 0. \end{cases}$$

Among these silting objects, $Z_0 = S_2 \oplus P_2$ and $Z_1 = S_2 \oplus S_1[1]$ are tilting objects.

6. HRS-TILTS AND RECOLLEMENTS

In this section we show that HRS-tilts of t-structures with respect to torsion theories ([15]) are *compatible* with the glueing of t-structures via recollements. Our notation is fixed as follows.

- $\mathcal{D}$ is a triangulated category admitting a recollement $\mathcal{R}$ of the form (2.1);
- $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ are bounded t-structures in $\mathcal{Y}$ and $\mathcal{X}$ respectively;
- $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is glued from $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ (theorem 2.2).

In fact (see [6], [21]), $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is glued with respect to $\mathcal{R}$ if and only if $j_! j^*(\mathcal{D}^{\leq 0}) \subseteq \mathcal{D}^{\leq 0}$ (or, equivalently, if and only if $j_* j^*(\mathcal{D}^{\geq 0}) \subseteq \mathcal{D}^{\geq 0}$) in which case the *restrictions* to $\mathcal{Y}$ and $\mathcal{X}$ satisfy

$$\begin{aligned} \mathcal{Y}^{\leq 0} &= i^*(\mathcal{D}^{\leq 0}), & \mathcal{Y}^{\geq 0} &= i^!(\mathcal{D}^{\geq 0}), \\ \mathcal{X}^{\leq 0} &= j^*(\mathcal{D}^{\leq 0}), & \mathcal{X}^{\geq 0} &= j^*(\mathcal{D}^{\geq 0}). \end{aligned}$$

Let $\mathcal{A}_{\mathcal{D}}$, $\mathcal{A}_{\mathcal{Y}}$ and $\mathcal{A}_{\mathcal{X}}$ be the hearts of these t-structures. In [6, section 1.4], it is shown that there is a *recollement of abelian categories* at the level of hearts

$$(6.1) \quad \begin{array}{ccccc} & \xleftarrow{p_i^*} & & \xleftarrow{p_j^!} & \\ \mathcal{A}_{\mathcal{Y}} & \xrightarrow{p_{i*}} & \mathcal{A}_{\mathcal{D}} & \xrightarrow{p_{j*}} & \mathcal{A}_{\mathcal{X}} \\ & \xleftarrow{p_i^!} & & \xleftarrow{p_{j*}} & \end{array}$$

We describe these functors explicitly. Let $\varepsilon_{\mathcal{D}} : \mathcal{A}_{\mathcal{D}} \hookrightarrow \mathcal{D}$ denote the full embedding (similarly $\varepsilon_{\mathcal{Y}} : \mathcal{A}_{\mathcal{Y}} \hookrightarrow \mathcal{Y}$ and $\varepsilon_{\mathcal{X}} : \mathcal{A}_{\mathcal{X}} \hookrightarrow \mathcal{X}$) and let $H_{\mathcal{D}}^i : \mathcal{D} \rightarrow \mathcal{A}_{\mathcal{D}}$, $i \in \mathbb{Z}$, denote the cohomological functors

with respect to the fixed t-structure in $\mathcal{D}$ (similarly $H_{\mathcal{Y}}^i$ and $H_{\mathcal{X}}^i$ for the t-structures in $\mathcal{Y}$ and $\mathcal{X}$). Then the functors in the recollement (6.1) are given by

$$\begin{aligned} {}^p i_* &= H_{\mathcal{Y}}^0 \circ i^* \circ \varepsilon_{\mathcal{D}}, \quad {}^p i^! = H_{\mathcal{Y}}^0 \circ i^! \circ \varepsilon_{\mathcal{D}}, \quad {}^p i_* = H_{\mathcal{D}}^0 \circ i_* \circ \varepsilon_{\mathcal{Y}}, \\ {}^p j_! &= H_{\mathcal{D}}^0 \circ j_! \circ \varepsilon_{\mathcal{X}}, \quad {}^p j_* = H_{\mathcal{D}}^0 \circ j_* \circ \varepsilon_{\mathcal{X}}, \quad {}^p j^* = H_{\mathcal{X}}^0 \circ j^* \circ \varepsilon_{\mathcal{D}}. \end{aligned}$$

See [14] for more on recollements of abelian categories.

Remark 6.1. Since for our fixed t-structures, i_* and j^* are t-exact (see, for example, [21] for details), we have that ${}^p i_* = i_* \circ \varepsilon_{\mathcal{Y}}$ and ${}^p j^* = j^* \circ \varepsilon_{\mathcal{D}}$. As a consequence, these two functors are exact.

Moreover, for any object A in $\mathcal{A}_{\mathcal{D}}$ there are exact sequences

$$(6.2) \quad 0 \rightarrow {}^p i_* H_{\mathcal{Y}}^{-1} {}^p i^* A \rightarrow {}^p j_! {}^p j^* A \rightarrow A \rightarrow {}^p i_* {}^p i^* A \rightarrow 0,$$

$$(6.3) \quad 0 \rightarrow {}^p i_* {}^p i^! A \rightarrow A \rightarrow {}^p j_* {}^p j^* A \rightarrow {}^p i_* H_{\mathcal{Y}}^1 {}^p i^* A \rightarrow 0.$$

Torsion pairs in $\mathcal{A}_{\mathcal{D}}$ satisfying certain conditions can be *restricted* to torsion pairs in $\mathcal{A}_{\mathcal{Y}}$ and $\mathcal{A}_{\mathcal{X}}$.

Lemma 6.2. *Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{A}_{\mathcal{D}}$. Then*

- (1) $({}^p i^*(\mathcal{T}), {}^p i^!(\mathcal{F}))$ is a torsion pair in $\mathcal{A}_{\mathcal{Y}}$;
- (2) *the following are equivalent:*
 - (i) $({}^p j^*(\mathcal{T}), {}^p j^*(\mathcal{F}))$ is a torsion pair in $\mathcal{A}_{\mathcal{X}}$,
 - (ii) ${}^p j_! {}^p j^*(\mathcal{T}) \subseteq \mathcal{T}$,
 - (iii) ${}^p j_* {}^p j^*(\mathcal{F}) \subseteq \mathcal{F}$.

Proof. (1): We first check the orthogonality condition for $({}^p i^*(\mathcal{T}), {}^p i^!(\mathcal{F}))$. We have that

$$\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^p i^*(\mathcal{T}), {}^p i^!(\mathcal{F})) = \mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^p i_* {}^p i^*(\mathcal{T}), {}^p i_* {}^p i^!(\mathcal{F}))$$

since ${}^p i_*$ is fully faithful. Consider the exact sequence (6.2) with $A \in \mathcal{T}$ and apply to it the functor $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}(-, B)$ for some $B \in \mathcal{F}$. Since $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}(A, B) = 0$, we conclude that $\mathrm{Hom}({}^p i_* {}^p i^*(A), B) = 0$, i.e., ${}^p i_* {}^p i^*(\mathcal{T}) \subseteq \mathcal{T}$. Similarly ${}^p i_* {}^p i^!(\mathcal{F}) \subseteq \mathcal{F}$ and, thus, $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^p i_* {}^p i^*(\mathcal{T}), {}^p i_* {}^p i^!(\mathcal{F})) = 0$.

Secondly, we produce a suitable short exact sequence for any object in $\mathcal{A}_{\mathcal{D}}$. By [6, remark 1.4.17.1], ${}^p i_*(\mathcal{A}_{\mathcal{Y}})$ is a Serre subcategory of $\mathcal{A}_{\mathcal{D}}$ and, thus, for a short exact sequence in $\mathcal{D}$,

$$0 \rightarrow K \rightarrow M \rightarrow L \rightarrow 0,$$

M lies in the ${}^p i_*(\mathcal{A}_{\mathcal{Y}})$ if and only if so do K and L . Let $M \in \mathcal{A}_{\mathcal{Y}}$ and consider a short exact sequence

$$0 \rightarrow \widetilde{M}_1 \rightarrow {}^p i_*(M) \rightarrow \widetilde{M}_2 \rightarrow 0$$

in $\mathcal{A}_{\mathcal{D}}$ with $\widetilde{M}_1 \in \mathcal{T}$ and $\widetilde{M}_2 \in \mathcal{F}$ (which exists since $(\mathcal{T}, \mathcal{F})$ is a torsion pair in $\mathcal{D}$). Since ${}^p i_*(\mathcal{A}_{\mathcal{Y}})$ is a Serre subcategory of $\mathcal{A}_{\mathcal{D}}$, there exist $M_1, M_2 \in \mathcal{A}_{\mathcal{Y}}$ such that $\widetilde{M}_i = {}^p i_*(M_i)$ for $i = 1, 2$. Now, ${}^p i_*$ is an exact full embedding (see remark 6.1) and, thus, we get a short exact sequence in $\mathcal{A}_{\mathcal{Y}}$

$$0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow 0,$$

with $M_1 \cong {}^p i^* {}^p i_*(M_1) = {}^p i^*(\widetilde{M}_1) \in {}^p i^*(\mathcal{T})$ and $M_2 \cong {}^p i^! {}^p i_*(M_2) = {}^p i^!(\widetilde{M}_2) \in {}^p i^!(\mathcal{F})$.

(2): By remark 6.1, ${}^p j^* : \mathcal{A}_{\mathcal{D}} \rightarrow \mathcal{A}_{\mathcal{X}}$ is exact. Therefore, $({}^p j^*(\mathcal{T}), {}^p j^*(\mathcal{F}))$ is a torsion pair in $\mathcal{A}_{\mathcal{X}}$ if and only if the orthogonality of the torsion pair is preserved, i.e. $\mathrm{Hom}_{\mathcal{A}_{\mathcal{X}}}({}^p j^*(\mathcal{T}), {}^p j^*(\mathcal{F})) = 0$. Using the adjunctions $({}^p j_!, {}^p j^*)$ and $({}^p j^*, {}^p j_*)$, this holds if and only if $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}({}^p j_! {}^p j^*(\mathcal{T}), \mathcal{F}) = 0$ (equivalent to (ii)) and if and only if $\mathrm{Hom}_{\mathcal{A}_{\mathcal{D}}}(\mathcal{T}, {}^p j_* {}^p j^*(\mathcal{F})) = 0$ (equivalent to (iii)). □

When the conditions of part (2) of the lemma are fulfilled, we say that $(\mathcal{T}, \mathcal{F})$ is *compatible* with the recollement (6.1). We will see that this compatibility condition is the exact requirement on a torsion pair so that the corresponding HRS-tilt of a glued t-structure is also obtained by glueing.

Proposition 6.3. *A torsion pair $(\mathcal{T}, \mathcal{F})$ in $\mathcal{A}_{\mathcal{D}}$ is compatible with respect to (6.1) if and only if the corresponding HRS-tilt of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is a t-structure glued with respect to the recollement (2.1).*

Proof. Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{A}_{\mathcal{D}}$. Denote by $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ the corresponding HRS-tilt, i.e., using the cohomological functors $H^i, i \in \mathbb{Z}$, of the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$,

$$\begin{aligned}\tilde{\mathcal{D}}^{\leq 0} &= \{X \in \mathcal{D} \mid H^0(X) \in \mathcal{T}, H^i(X) = 0, \forall i > 0\}, \\ \tilde{\mathcal{D}}^{\geq 0} &= \{X \in \mathcal{D} \mid H^{-1}(X) \in \mathcal{F}, H^i(X) = 0, \forall i < -1\}.\end{aligned}$$

By [6], the HRS-tilt is glued if and only if $j_! j^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq \tilde{\mathcal{D}}^{\leq 0}$. Denote by $\tau^{\leq -1}$ the truncation associated to $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$. For any $X \in \tilde{\mathcal{D}}^{\leq 0}$, there is a triangle in $\mathcal{D}$ of the form

$$\tau^{\leq -1} X \rightarrow X \rightarrow H^0 X \rightarrow (\tau^{-1} X)[1],$$

since, by definition, $\tilde{\mathcal{D}}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$. Applying to it the functor $j_! j^*$, we get another triangle

$$j_! j^*(\tau^{\leq -1} X) \rightarrow j_! j^*(X) \rightarrow j_! j^*(H^0 X) \rightarrow j_! j^*(\tau^{\leq -1} X)[1].$$

Since $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ is glued, $j_! j^*(\mathcal{D}^{\leq -1}) \subseteq \mathcal{D}^{\leq -1} \subseteq \tilde{\mathcal{D}}^{\leq 0}$. Hence, $j_! j^*(\tau^{\leq -1} X) \in \tilde{\mathcal{D}}^{\leq 0}$ and, thus, $j_! j^*(X) \in \tilde{\mathcal{D}}^{\leq 0}$ if and only if $j_! j^*(H^0 X) \in \tilde{\mathcal{D}}^{\leq 0}$. Since X is arbitrary, this means precisely that $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ can be restricted if and only if the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with (6.1). $\square$

The question that naturally follows is whether, under the compatibility condition on the torsion pair in $\mathcal{D}$, the restrictions of the HRS-tilt of a glued t-structure in $\mathcal{D}$ are precisely the HRS-tilts of the restricted t-structures on $\mathcal{Y}$ and $\mathcal{X}$ with respect to the restricted torsion pairs.

Theorem 6.4. *Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair in $\mathcal{A}_{\mathcal{D}}$, compatible with (6.1) and with restrictions $(\mathcal{T}_{\mathcal{Y}}, \mathcal{F}_{\mathcal{Y}})$ and $(\mathcal{T}_{\mathcal{X}}, \mathcal{F}_{\mathcal{X}})$ in $\mathcal{A}_{\mathcal{Y}}$ and $\mathcal{A}_{\mathcal{X}}$ respectively. Let $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ and $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ be the corresponding HRS-tilts of $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ respectively. Then the HRS-tilt $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ is obtained by glueing $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ and $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ with respect to the recollement (2.1).*

Proof. Let $H_{\mathcal{D}}^i, H_{\mathcal{X}}^i$ and $H_{\mathcal{Y}}^i, i \in \mathbb{Z}$, denote the cohomological functors associated with $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ respectively. We have to show that the following holds:

- (i) $j^*(\tilde{\mathcal{D}}^{\leq 0}) = \tilde{\mathcal{X}}^{\leq 0}, j^*(\tilde{\mathcal{D}}^{\geq 0}) = \tilde{\mathcal{X}}^{\geq 0};$
- (ii) $i^*(\tilde{\mathcal{D}}^{\leq 0}) = \tilde{\mathcal{Y}}^{\leq 0}, i^*(\tilde{\mathcal{D}}^{\geq 0}) = \tilde{\mathcal{Y}}^{\geq 0}.$

By definition, we have

$$j^*(\tilde{\mathcal{D}}^{\leq 0}) = \{j^*(X) \mid X \in \mathcal{D}, H_{\mathcal{D}}^0(X) \in \mathcal{T}, H_{\mathcal{D}}^i(X) = 0, \forall i > 0\},$$

and, since j^* is t-exact ([6]), $H_{\mathcal{X}}^i(j^*X) = j^*(H_{\mathcal{D}}^i(X))$, thus showing that

$$j^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq \{j^*(X) \mid X \in \mathcal{D}, H_{\mathcal{X}}^0(j^*X) \in j^*\mathcal{T}, H_{\mathcal{X}}^i(j^*X) = 0, \forall i > 0\} = \tilde{\mathcal{X}}^{\leq 0}.$$

Similarly, we get that

$$j^*(\tilde{\mathcal{D}}^{\geq 0}) \subseteq \{j^*(X) \mid X \in \mathcal{D}, H_{\mathcal{X}}^{-1}(j^*X) \in j^*\mathcal{T}, H_{\mathcal{X}}^i(j^*X) = 0, \forall i < -1\} = \tilde{\mathcal{X}}^{\geq 0}.$$

Since $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement, by proposition 6.3, the HRS-tilt $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ can be restricted and, thus, $(j^*(\tilde{\mathcal{D}}^{\leq 0}), j^*(\tilde{\mathcal{D}}^{\geq 0}))$ is a t-structure in $\mathcal{X}$. Since $j^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq \tilde{\mathcal{X}}^{\leq 0}$, we also have that $j^*(\tilde{\mathcal{D}}^{\geq 0}) \supseteq \tilde{\mathcal{X}}^{\geq 0}$ and, thus, $j^*(\tilde{\mathcal{D}}^{\geq 0}) = \tilde{\mathcal{X}}^{\geq 0}$. This proves (i).

To prove (ii), observe that i^* is right t-exact (i.e., $i^*(\mathcal{D}^{\leq 0}) \subseteq \mathcal{Y}^{\leq 0}$) and that

$$i^*(\tilde{\mathcal{D}}^{\leq 0}) = \{i^*(X) \mid X \in \mathcal{D}, H_{\mathcal{D}}^0(X) \in \mathcal{T}, H_{\mathcal{D}}^i(X) = 0, \forall i > 0\}.$$

By [6] (proposition 1.3.17 (ii)), for $X \in \tilde{\mathcal{D}}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$, $H_{\mathcal{Y}}^0(i^*(X)) = {}^p i^* H_{\mathcal{D}}^0(X)$ which belongs to ${}^p i^*(\mathcal{T})$. Since $i^*(\tilde{\mathcal{D}}^{\leq 0}) \subset i^*(\mathcal{D}^{\leq 0}) \subseteq \mathcal{Y}^{\leq 0}$, we get that $H_{\mathcal{Y}}^i(i^*(\tilde{\mathcal{D}}^{\leq 0})) = 0$, for all $i > 0$, and thus

$$i^*(\tilde{\mathcal{D}}^{\leq 0}) \subseteq \{i^*(X) \mid X \in \mathcal{D}, H_{\mathcal{Y}}^0(i^*X) \in {}^p i^*\mathcal{T}, H_{\mathcal{Y}}^i(i^*X) = 0, \forall i > 0\} = \tilde{\mathcal{Y}}^{\leq 0}.$$

Analogously, we may conclude that

$$i^*(\tilde{\mathcal{D}}^{\geq 0}) \subseteq \{i^*(X) \mid X \in \mathcal{D}, H_{\mathcal{Y}}^{-1}(i^*X) \in {}^p i^*\mathcal{F}, H_{\mathcal{Y}}^i(i^*X) = 0, \forall i < -1\} = \tilde{\mathcal{Y}}^{\geq 0}.$$

The same argument as before shows that an equality holds, as wanted. $\square$

The previous result showed that an HRS-tilt with respect to a compatible torsion pair is glued. We may now ask whether HRS-tilts on the sides of the recollement (2.1) glue to an HRS-tilt in the middle. The following proposition answers this question positively.

Remark 6.5. HRS-tilts span an important class of t-structures. In fact, it is known from [7, theorem 3.1] and [29, proposition 2.1] that, for a bounded t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, the bounded t-structures $(\mathcal{C}^{\leq 0}, \mathcal{C}^{\geq 0})$ such that $\mathcal{D}^{\leq -1} \subseteq \mathcal{C}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$ are precisely those obtained from $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ by an HRS-tilt with respect to a torsion pair.

Proposition 6.6. *Let $(\mathcal{T}_{\mathcal{Y}}, \mathcal{F}_{\mathcal{Y}})$ and $(\mathcal{T}_{\mathcal{X}}, \mathcal{F}_{\mathcal{X}})$ be torsion pairs in $\mathcal{A}_{\mathcal{Y}}$ and $\mathcal{A}_{\mathcal{X}}$ respectively. Then $(\mathcal{T}, \mathcal{F})$ defined by*

$$\begin{aligned} \mathcal{T} &= \{A \in \mathcal{A}_{\mathcal{D}} \mid {}^p i^* A \in \mathcal{T}_{\mathcal{Y}}, {}^p j^* A \in \mathcal{T}_{\mathcal{X}}\} \\ \mathcal{F} &= \{A \in \mathcal{A}_{\mathcal{D}} \mid {}^p i^! A \in \mathcal{F}_{\mathcal{Y}}, {}^p j^* A \in \mathcal{F}_{\mathcal{X}}\}. \end{aligned}$$

is a torsion pair in $\mathcal{A}_{\mathcal{D}}$, compatible with (6.1), whose restrictions are given by $(\mathcal{T}_{\mathcal{X}}, \mathcal{F}_{\mathcal{X}})$ and $(\mathcal{T}_{\mathcal{Y}}, \mathcal{F}_{\mathcal{Y}})$, and the HRS-tilt of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to $(\mathcal{T}, \mathcal{F})$ is the glueing of the HRS-tilts of $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ with respect to $(\mathcal{T}_{\mathcal{Y}}, \mathcal{F}_{\mathcal{Y}})$ and $(\mathcal{T}_{\mathcal{X}}, \mathcal{F}_{\mathcal{X}})$ respectively.

Proof. We will use remark 6.5 to show that $(\mathcal{T}, \mathcal{F})$ is a torsion pair, i.e., we will show that the pair $(\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$ formally defined as the HRS-tilt of $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$ with respect to $(\mathcal{T}, \mathcal{F})$ is a t-structure (satisfying $\mathcal{D}^{\leq -1} \subseteq \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$). Denote by $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ and $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ the HRS-tilts of $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$ and $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ at $(\mathcal{T}_{\mathcal{Y}}, \mathcal{F}_{\mathcal{Y}})$ and $(\mathcal{T}_{\mathcal{X}}, \mathcal{F}_{\mathcal{X}})$ respectively and let $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0})$ be the glueing of $(\tilde{\mathcal{Y}}^{\leq 0}, \tilde{\mathcal{Y}}^{\geq 0})$ and $(\tilde{\mathcal{X}}^{\leq 0}, \tilde{\mathcal{X}}^{\geq 0})$ with respect to the fixed recollement, i.e.,

$$\begin{aligned} \tilde{\mathcal{D}}^{\leq 0} &= \{X \in \mathcal{D} \mid i^*(X) \in \tilde{\mathcal{Y}}^{\leq 0}, j^*(X) \in \tilde{\mathcal{X}}^{\leq 0}\} \\ \tilde{\mathcal{D}}^{\geq 0} &= \{X \in \mathcal{D} \mid i^!(X) \in \tilde{\mathcal{Y}}^{\geq 0}, j^*(X) \in \tilde{\mathcal{X}}^{\geq 0}\}. \end{aligned}$$

We will show that $(\tilde{\mathcal{D}}^{\leq 0}, \tilde{\mathcal{D}}^{\geq 0}) = (\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}, \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\geq 0})$. As usual, $H_{\mathcal{D}}^i, H_{\mathcal{X}}^i$ and $H_{\mathcal{Y}}^i$, $i \in \mathbb{Z}$, denote the cohomological functors associated with $(\mathcal{D}^{\leq 0}, \mathcal{D}^{\geq 0})$, $(\mathcal{X}^{\leq 0}, \mathcal{X}^{\geq 0})$ and $(\mathcal{Y}^{\leq 0}, \mathcal{Y}^{\geq 0})$, respectively.

First we show that $\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}$ is a subcategory of $\tilde{\mathcal{D}}^{\leq 0}$. Take X in $\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0} (\subseteq \mathcal{D}^{\leq 0})$. Since i^* is right t-exact, $i^*(X)$ lies in $\mathcal{Y}^{\leq 0}$, i.e., $H_{\mathcal{Y}}^i(i^*X) = 0$, for all $i > 0$. By [6] (proposition 1.3.17(ii)), $H_{\mathcal{Y}}^0(i^*X) = i^*H_{\mathcal{D}}^0(X)$ lies in $i^*(\mathcal{T}) \subseteq \mathcal{T}_{\mathcal{Y}}$, showing that $i^*(X) \in \tilde{\mathcal{Y}}^{\leq 0}$. Since j^* is t-exact, we have $H_{\mathcal{X}}^i(j^*X) = j^*H_{\mathcal{D}}^i(X) = 0$, for all $i > 0$, and $H_{\mathcal{X}}^0(j^*X) = j^*H_{\mathcal{D}}^0(X)$ lies in $j^*(\mathcal{T}) \subseteq \mathcal{T}_{\mathcal{X}}$, i.e., $j^*X \in \tilde{\mathcal{X}}^{\leq 0}$. Hence $X \in \tilde{\mathcal{D}}^{\leq 0}$.

Now we prove that $\tilde{\mathcal{D}}^{\leq 0}$ is a subcategory of $\mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}$. Take X in $\tilde{\mathcal{D}}^{\leq 0}$. By definition of HRS-tilt, we have that $\tilde{\mathcal{Y}}^{\leq 0} \subset \mathcal{Y}^{\leq 0}$ and $\tilde{\mathcal{X}}^{\leq 0} \subseteq \mathcal{X}^{\leq 0}$. This implies that $\tilde{\mathcal{D}}^{\leq 0} \subseteq \mathcal{D}^{\leq 0}$ and, hence, $H_{\mathcal{D}}^i(X) = 0$, for all $i > 0$. By [6] (proposition 1.3.17(ii)), we have ${}^p i^* H_{\mathcal{D}}^0(X) = H_{\mathcal{Y}}^0(i^*X)$ which belongs to $\mathcal{T}_{\mathcal{Y}}$. By the t-exactness of j^* , ${}^p j^* H_{\mathcal{D}}^0(X) = H_{\mathcal{X}}^0(j^*X)$ which, thus, belongs to $\mathcal{T}_{\mathcal{X}}$. This means that, by definition of $\mathcal{T}$, $H_{\mathcal{D}}^0(X)$ belongs to $\mathcal{T}$. Thus X belongs to $\tilde{\mathcal{D}}^{\leq 0}$ and $\tilde{\mathcal{D}}^{\leq 0} = \mathcal{D}_{(\mathcal{T}, \mathcal{F})}^{\leq 0}$, as wanted.

It remains to show that $(\mathcal{T}, \mathcal{F})$ is compatible and that the restrictions are as expected. By lemma 6.2, we need to show ${}^p j_! {}^p j^* \mathcal{T} \subseteq \mathcal{T}$. This is true since, for any $A \in \mathcal{T}$, ${}^p i^* {}^p j_! {}^p j^* A = 0$ and ${}^p j^* {}^p j_! {}^p j^* A = {}^p j^* A \subseteq \mathcal{T}_{\mathcal{X}}$. It follows that $({}^p i^*(\mathcal{T}), {}^p i^!(\mathcal{F}))$ and $({}^p j^*(\mathcal{T}), {}^p j^*(\mathcal{F}))$ are torsion pairs in $\mathcal{Y}$ and $\mathcal{X}$ respectively. On the other hand, by definition of $(\mathcal{T}, \mathcal{F})$ we have

$${}^p i^*(\mathcal{T}) \subseteq \mathcal{T}_{\mathcal{Y}}, \quad {}^p i^!(\mathcal{F}) \subseteq \mathcal{F}_{\mathcal{Y}},$$

$${}^p j^*(\mathcal{T}) \subseteq \mathcal{T}_{\mathcal{X}}, \quad {}^p j^*(\mathcal{F}) \subseteq \mathcal{F}_{\mathcal{X}},$$

and, thus, the inclusions are actually equalities, as wanted. $\square$

By subsection 2.4, we know that irreducible silting mutations correspond to HRS-tilts (see theorem 2.10) in which either the torsion class or the torsion-free class are given by a simple object in the heart. We analyse such torsion pairs which are compatible with (6.1), which we will call *mutation torsion pairs*.

Remark 6.7. Recall that there is a functor $j_{!*} : \mathcal{A}_{\mathcal{X}} \rightarrow \mathcal{A}_{\mathcal{D}}$, called the intermediate image functor, such that every simple object in $\mathcal{A}_{\mathcal{D}}$ is either of the form $i_* S$ for some simple object in $\mathcal{A}_{\mathcal{Y}}$ or of the form $j_{!*} S$ for some simple object in $\mathcal{A}_{\mathcal{X}}$. For details, check [6, section 1.4].

Proposition 6.8. *Let $(\mathcal{T}, \mathcal{F})$ be a mutation torsion pair in $\mathcal{A}_{\mathcal{D}}$. Then $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement (6.1) if and only if one of the following holds.*

- (1) $\mathcal{T} = \text{add}(i_* S_Y)$ for some simple object S_Y in $\mathcal{A}_{\mathcal{Y}}$;
- (2) $\mathcal{F} = \text{add}(i_* S_Y)$ for some simple object S_Y in $\mathcal{A}_{\mathcal{Y}}$;
- (3) there is a simple object $S_X \in \mathcal{A}_{\mathcal{X}}$ such that ${}^p j_! S_X$ is simple and $\mathcal{T} = \text{add}({}^p j_! S_X)$;
- (4) there is a simple object $S_X \in \mathcal{A}_{\mathcal{X}}$ such that ${}^p j_* S_X$ is simple and $\mathcal{F} = \text{add}({}^p j_* S_X)$.

Moreover, in each case, the restrictions are a trivial torsion pair on one side and a mutation torsion pair on the other.

Proof. This can be showed using lemma 6.2 on a case by case analysis of the possible mutation torsion pairs in $\mathcal{A}_{\mathcal{D}}$. Using remark 6.7 we get the following four cases, where S_Y is a simple object in $\mathcal{A}_{\mathcal{Y}}$ and S_X is a simple object in $\mathcal{A}_{\mathcal{X}}$.

- (1) Suppose that $\mathcal{T} = \text{add}(i_* S_Y)$. Since $j^* i_* = 0$, we have that ${}^p j_! j^* \mathcal{T} = 0$ and, thus, the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement. The restriction to $\mathcal{X}$ is $(0, \mathcal{X})$ and the restriction to $\mathcal{Y}$ is the mutation torsion pair with torsion class $\mathcal{T}_{\mathcal{Y}} = \text{add}(S_Y)$.

- (2) Suppose that $\mathcal{F} = \text{add}(i_*S_Y)$. Since $j^*i_* = 0$, we have that ${}^Pj_*j^*\mathcal{T} = 0$ and, thus, the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement. The restriction to $\mathcal{X}$ is $(\mathcal{X}, 0)$ and the restriction to $\mathcal{Y}$ is the mutation torsion pair with torsion-free class $\mathcal{F}_{\mathcal{Y}} = \text{add}(S_Y)$.
- (3) Suppose that $\mathcal{T} = \text{add}(j_{!*}S_X)$. By lemma 6.2 the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement if and only if ${}^Pj_!j^*(j_{!*}S_X) \in \text{add}(j_{!*}S_X)$. Since $j^*j_{!*}$ is naturally equivalent to the identity functor, this amounts to ${}^Pj_!S_X \in \text{add}(j_{!*}S_X)$, which is equivalent to ${}^Pj_!S_X \cong j_{!*}S_X$ because $j^*{}^Pj_!$ is also naturally equivalent to the identity functor. The last condition holds if and only if ${}^Pj_!S_X$ is simple. This follows from the fact that there is always a natural epimorphism ${}^Pj_!S_X \rightarrow j_{!*}S_X$. The restriction to $\mathcal{Y}$ is the pair $(0, \mathcal{Y})$ and the restriction to $\mathcal{X}$ is the mutation torsion pair given by the torsion class $\mathcal{T}_{\mathcal{X}} = \text{add}(S_X)$.
- (4) Suppose that $\mathcal{F} = \text{add}(j_{!*}S_X)$. By lemma 6.2 the torsion pair $(\mathcal{T}, \mathcal{F})$ is compatible with the recollement if and only if ${}^Pj_*j^*(j_{!*}S_X) \in \text{add}(j_{!*}S_X)$. Since $j^*j_{!*}$ is naturally equivalent to the identity functor, this amounts to ${}^Pj_*S_X \in \text{add}(j_{!*}S_X)$, which is equivalent to ${}^Pj_*S_X \cong j_{!*}S_X$ because $j^*{}^Pj_*$ is also naturally equivalent to the identity functor. The last condition holds if and only if ${}^Pj_*S_X$ is simple. This follows from the fact that there is always a natural monomorphism $j_{!*}S_X \rightarrow {}^Pj_*S_X$. The restriction to $\mathcal{Y}$ is the pair $(\mathcal{Y}, 0)$ and the restriction to $\mathcal{X}$ is the mutation torsion pair given by the torsion-free class $\mathcal{F}_{\mathcal{X}} = \text{add}(S_X)$.

□

Suppose now that $\mathcal{D}$ has a Serre functor. We finish this section with an observation on the compatibility of irreducible silting mutation with glueing (via the recollement $\mathcal{R}_{\mathcal{U}}$, since our focus is on t-structures rather than co-t-structures). Note that, in general, items (3) and (4) of the proposition above will not occur. Therefore, translating items (1) and (2) of the proposition in terms of irreducible silting mutation, we get the following corollary, also appearing in [1] (lemma 2.40).

Corollary 6.9. *Let Y and X be the silting objects associated to the fixed bounded t-structures in $\mathcal{Y}$ and $\mathcal{X}$, respectively. Suppose Y' is an indecomposable direct summand of Y . Let Z be the glued silting object with respect to the recollement $\mathcal{R}_{\mathcal{U}}$ (i.e., compatible with the glueing of t-structures via $\mathcal{R}_{\mathcal{U}}$) of Y and X . Then there is an indecomposable direct summand Z' of K_Y such that the glued silting (with respect to $\mathcal{R}_{\mathcal{U}}$) of $\mu_{Y'}^+Y$ (respectively, $\mu_{Y'}^-Y$) and X is precisely $\mu_{Z'}^+Z$ (respectively, $\mu_{Z'}^-Z$).*

REFERENCES

1. Takuma Aihara and Osamu Iyama, *Silting mutation in triangulated categories*, J. Lon. Math. Soc. **85** (2012), no. 3, 633–668.
2. Leovigildo Alonso Tarrío, Ana Jeremías López, and María José Souto Salorio, *Construction of t-structures and equivalences of derived categories*, Trans. Amer. Math. Soc. **355** (2003), no. 6, 2523–2543 (electronic).
3. Lidia Angeleri Hügel, Steffen Koenig, and Qunhua Liu, *Recollements and tilting objects*, J. Pure Appl. Algebra **215** (2011), no. 4, 420–438.
4. ———, *Jordan-Hölder theorems for derived module categories of piecewise hereditary algebras*, J. Algebra **352** (2012), 361–381.
5. ———, *On the uniqueness of stratifications of derived module categories*, J. Algebra **359** (2012), 120–137.
6. Alexander A. Beilinson, Joseph Bernstein, and Pierre Deligne, *Analyse et topologie sur les espaces singuliers*, Astérisque, vol. 100, Soc. Math. France, 1982 (French).
7. Apostolos Beligiannis and Idun Reiten, *Homological and homotopical aspects of torsion theories*, Mem. Amer. Math. Soc. **188** (2007), no. 883, viii+207.
8. A. I. Bondal and M. M. Kapranov, *Representable functors, Serre functors, and reconstructions*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1183–1205, 1337.
9. M. V. Bondarko, *Weight structures vs. t-structures; weight filtrations, spectral sequences, and complexes (for motives and in general)*, J. K-Theory **6** (2010), no. 3, 387–504.

10. Tom Bridgeland, *t-structures on some local Calabi-Yau varieties*, J. Algebra **289** (2005), no. 2, 453–483.
11. Aslak Bakke Buan, Idun Reiten, and Hugh Thomas, *Three kinds of mutation*, J. Algebra **339** (2011), 97–113.
12. Edward Cline, Brian Parshall and Leonard L. Scott, *Algebraic stratification in representation categories*, J. Algebra **117** (1988), no. 2, 504–521.
13. E. Cline, B. Parshall, L. Scott, *Stratifying endomorphism algebras*, Mem. Amer. Math. Soc. **124** (1996), no. 591.
14. Vincent Franjou and Teimuraz Pirashvili, *Comparison of abelian categories recollements*, Doc. Math. **9** (2004), 41–56 (electronic).
15. Dieter Happel, Idun Reiten, and Sverre O. Smalø, *Tilting in abelian categories and quasitilted algebras*, Mem. Amer. Math. Soc. **120** (1996), no. 575, viii+ 88.
16. Peter Jørgensen, *Reflecting recollements*, Osaka J. Math. **47** (2010), no. 1, 209–213.
17. Bernhard Keller and Pedro Nicolás, *Cluster hearts and cluster tilting objects*, in preparation.
18. Bernhard Keller and Dieter Vossieck, *Aisles in derived categories*, Bull. Soc. Math. Belg. Sér. A **40** (1988), no. 2, 239–253.
19. Steffen Koenig and Dong Yang, *Silting objects, simple-minded collections, t-structures and co-t-structures for finite-dimensional algebras*, arXiv:1203.5657.
20. Sefi Ladkani, *Derived equivalences of triangular matrix rings arising from extensions of tilting modules*, Algebr. Represent. Theory **14** (2011), 57–74.
21. Qunhua Liu and Jorge Vitória, *t-structures via recollements for piecewise hereditary algebras*, J. Pure Appl. Algebra **216** (2012), no. 4, 837–849.
22. Octavio Mendoza, Edith C. Sáenz, Valente Santiago, and María José Souto Salorio, *Auslander-Buchweitz context and co-t-structures*, Appl. Categor. Struct., published online, doi:10.1007/s10485-011-9271-2, arXiv:1002.4604v3.
23. David Pauksztello, *Compact corigid objects in triangulated categories and co-t-structures*, Cent. Eur. J. Math. **6** (2008), no. 1, 25–42.
24. Idun Reiten and Michel Van den Bergh, *Noetherian hereditary abelian categories satisfying Serre duality*, J. Amer. Math. Soc. **15** (2002), no. 2, 295–366 (electronic).
25. Jeremy Rickard, *Morita theory for derived categories*, J. Lon. Math. Soc. **39** (1989), 436–456.
26. Jeremy Rickard and Aidan Schofield, *Cocovers and tilting modules*, Math. Proc. Cambridge Philos. Soc. **106** (1989), no. 1, 1–5.
27. María José Souto Salorio and Sonia Trepode, *T-structures on the Bounded Derived Category of the Kronecker Algebra*, Appl Categor Struct, in press, doi:10.1007/s10485-011-9248-1.
28. Zhongmei Wang, *Nondegeneration and boundedness of t-structure induced by recollement*, Xiamen Daxue Xuebao Ziran Kexue Ban **45** (2006), no. 1, 10–13.
29. Jonathan Woolf, *Stability conditions, torsion theories and tilting*, J. Lond. Math. Soc. (2) **82** (2010), no. 3, 663–682.

QUNHUA LIU, JORGE VITÓRIA, DONG YANG, INSTITUTE FOR ALGEBRA AND NUMBER THEORY, UNIVERSITY OF STUTTGART, PFAFFENWALDRING 57, D-70569 STUTTGART, GERMANY

E-mail address: qliu@mathematik.uni-stuttgart.de

E-mail address: vitoria@mathematik.uni-stuttgart.de

E-mail address: dongyang2002@gmail.com